

LOCUS

Definition: The set of all points (and only those points) which satisfy the given geometrical condition(s) (or properties) is called a locus.

Eg. The set of points in a plane which are at a constant distance r from a given point C is a locus. Here the locus is a circle.

2. The set of points in a plane which are equidistant from two given points A and B is a locus. Here the locus is a straight line and it is the perpendicular bisector of the line segment joining A and B .

EQUATION OF A LOCUS

An equation $f(x, y) = 0$ is said to be the equation of a locus S if every point of S satisfies $f(x, y) = 0$ and every point that satisfies $f(x, y) = 0$ belongs to S .

An equation of a locus is an algebraic description of the locus. This can be obtained in the following way

- (i) Consider a point $P(x, y)$ on the locus
- (ii) Write the geometric condition(s) to be satisfied by P in terms of an equation or in equation in symbols.
- (iii) Apply the proper formula of coordinate geometry and translate the geometric condition(s) into an algebraic equation.
- (iv) Simplify the equation so that it is free from radicals.

The equation thus obtained is the required equation of locus.


EXERCISE

I.

1. Find the equation of locus of a point which is at a distance 5 from A(4, -3).

Sol. Let P(x, y) be a point in the locus.

Given A(4, -3)



Given that CP = 5

$$\Rightarrow CP^2 = 25$$

$$\Rightarrow (x - 4)^2 + (y + 3)^2 = 25$$

$$\Rightarrow x^2 - 8x + 16 + y^2 + 6y + 9 - 25 = 0$$


\therefore Equation of the locus of P is:

$$x^2 + y^2 - 8x + 6y = 0$$

2. Find the equation of locus of a point which is equidistant from the points A(-3, 2) and B(0, 4).

Sol. Given points are A(-3, 2), B(0, 4)

Let P(x, y) be any point in the locus



Given that PA = PB

$$\Rightarrow PA^2 = PB^2$$

$$\Rightarrow (x + 3)^2 + (y - 2)^2 = (x - 0)^2 + (y - 4)^2$$


$$\Rightarrow x^2 + 6x + 9 + y^2 - 4y + 4 = x^2 + y^2 - 8y + 16$$

$$\Rightarrow 6x + 4y = 3 \text{ is the equation of the locus.}$$

3. Find the equation of locus of a point P such that the distance of P from the origin is twice the distance of P from A (1, 2).

Sol. Given points are O (0, 0), A (1, 2)

Let P(x, y) be any point in the locus



Given that $OP = 2AP$

$$\Rightarrow OP^2 = 4AP^2$$

$$\Rightarrow x^2 + y^2 = 4[(x-1)^2 + (y-2)^2]$$

$$= 4(x^2 - 2x + 1 + y^2 - 4y + 4)$$

$$\Rightarrow x^2 + y^2 = 4x^2 + 4y^2 - 8x - 16y + 20$$

\therefore Equation to the locus of P is


$$3x^2 + 3y^2 - 8x - 16y + 20 = 0$$

4. Find the equation of locus of a point which is equidistant from the coordinate axes.

Sol. Let $P(x, y)$ be any point in the locus.

Let PM = perpendicular distance of P from X-axis. = $|x|$

Let PN = perpendicular distance of P from Y-axis. = $|y|$



Given $PM = PN \Rightarrow |x| = |y|$

Squaring on both sides, $x^2 = y^2$

Therefore, Locus of P is $x^2 - y^2 = 0$

5. Find the equation of locus of a point equidistant from $A(2, 0)$ and the Y-axis.

Sol. Given point is $A(2, 0)$

Let $P(x, y)$ be any point in the locus.

Draw PN perpendicular to Y-axis. = $|x|$


Given that is $PA = PN$

$$\Rightarrow PA^2 = PN^2$$


$$\Rightarrow (x-2)^2 + (y-0)^2 = x^2$$

$$\Rightarrow x^2 - 4x + 4 + y^2 = x^2$$

$$\therefore \text{Locus of P is } y^2 - 4x + 4 = 0$$



6. Find the equation of locus of a point P, the square of whose distance from the origin is 4 times its y coordinates.



Sol. Let $P(x, y)$ be any point in the locus.

$$\text{Now } OP^2 = x^2 + y^2$$

Given condition is $OP^2 = 4y \Rightarrow x^2 + y^2 = 4y$
 Equation of the locus of P is $x^2 + y^2 - 4y = 0$

- 7. Find the equation of locus of a point P such that $PA^2 + PB^2 = 2c^2$, where $A = (a, 0)$, $B = (-a, 0)$ and $0 < |a| < |c|$.**

Sol. Let $P(x, y)$ be a point in locus.

Given $A = (a, 0)$, $B = (-a, 0)$

Given that $PA^2 + PB^2 = 2c^2$

$$\Rightarrow (x - a)^2 + (y - 0)^2 + (x + a)^2 + (y - 0)^2 = 2c^2$$

$$\Rightarrow x^2 - 2ax + a^2 + y^2 + x^2 + 2ax + a^2 + y^2 = 2c^2$$

$$\Rightarrow 2x^2 + 2y^2 = 2c^2 - 2a^2$$

$\therefore x^2 + y^2 = c^2 - a^2$ is the locus of P.

II.

- 1. Find the equation of locus of P, if the line segment joining $(2, 3)$ and $(-1, 5)$ subtends a right angle at P.**

Sol. Given points $A(2, 3)$, $B(-1, 5)$.

Let $P(x, y)$ be any point in the locus.

Given condition is: $\angle APB = 90^\circ$


$$\Rightarrow (\text{Slope of } \overline{AP}) (\text{Slope of } \overline{BP}) = -1$$

$$\Rightarrow \frac{y-3}{x-2} \cdot \frac{y-5}{x+1} = -1$$

$$(y-3)(y-5) + (x-2)(x+1) = 0$$

$$x^2 + y^2 - x - 8y + 13 = 0$$

\therefore Locus of P is $x^2 + y^2 - x - 8y + 13 = 0$



- 2. The ends of the hypotenuse of a right angled triangle are $(0, 6)$ and $(6, 0)$. Find the equation of locus of its third vertex.**

Sol. Same as above.

- 3. Find the equation of locus of a point, the difference of whose distances from $(-5, 0)$ and $(5, 0)$ is 8 units.**

Sol. Given points are $A(5, 0)$, $B(-5, 0)$

Let $P(x, y)$ be any point in the locus

Given $|PA - PB| = 8$

$$\Rightarrow PA - PB = \pm 8$$

$$\Rightarrow PA = \pm 8 + PB$$

Squaring on both sides

$$PA^2 = 64 + PB^2 \pm 16PB$$

$$PA^2 - 64 - PB^2 = \pm 16PB$$

$$\Rightarrow (x-5)^2 + y^2 - (x+5)^2 - y^2 - 64 = \pm 16PB$$

$$-4 \cdot 5 \cdot x - 64 = \pm 16PB$$

$$-5x - 16 = \pm 4PB$$

Squaring on both sides

$$\begin{aligned} 25x^2 + 256 + 160x &= 16(PB)^2 \\ &= 16[(x+5)^2 + y^2] \\ &= 16x^2 + 400 + 160x + 16y^2 \end{aligned}$$

$$\Rightarrow 9x^2 - 16y^2 = 144$$

$$\Rightarrow \frac{9x^2}{144} - \frac{16y^2}{144} = 1$$

$$\Rightarrow \text{locus of P is } \frac{x^2}{16} - \frac{y^2}{9} = 1$$

4. Find the equation of locus of P, if A(4, 0), B(-4, 0) and $|PA - PB| = 4$.

Sol. Same as above.

5. Find the equation of locus of a point, the sum of whose distances from (0, 2) and (0, -2) is 6.

Sol. Given points are A (0, 2) and B (0, -2)

Let P(x, y) be any point in the locus.

$$\text{Given } PA + PB = 6$$

$$\Rightarrow PA = 6 - PB$$

Squaring on both sides

$$PA^2 = 36 + PB^2 - 12PB$$

$$12PB = PB^2 - PA^2 + 36$$

$$= x^2 + (y+2)^2 - [x^2 + (y-2)^2] + 36$$

$$\Rightarrow 12PB = 4 \cdot 2 \cdot y + 36$$

$$\Rightarrow 3PB = 2y + 9$$

squaring on both sides

$$9PB^2 = 4y^2 + 36y + 81$$

$$\Rightarrow 9[x^2 + (y+2)^2] = 4y^2 + 36y + 81$$

$$\Rightarrow 9x^2 + 9y^2 + 36 + 36y = 4y^2 + 36y + 81$$

$$\Rightarrow 9x^2 + 5y^2 = 45$$

$$\Rightarrow \frac{9x^2}{45} + \frac{5y^2}{45} = 1 \Rightarrow \text{Locus of P is } \frac{x^2}{5} + \frac{y^2}{9} = 1.$$

6. Find the equation of locus of P, if A(2, 3), B(2, -3) and $PA + PB = 8$.

Sol. Same as above.

7. **A(5, 3) and B(3, -2) are two fixed points. Find the equation of locus of P, so that the area of triangle PAB is 9.**

Sol. Given points are A(5, 3), B(3, -2)

Let P(x, y) be a point in the locus.

Given, area of $\Delta APB = 9$.

$$\Rightarrow \frac{1}{2} \begin{vmatrix} x-5 & y-3 \\ 3-5 & -2-3 \end{vmatrix} = 9$$

$$\Rightarrow \begin{vmatrix} x-5 & y-3 \\ -2 & -5 \end{vmatrix} = 18$$

$$\Rightarrow |-5x + 25 + 2y - 6| = 18$$

$$\Rightarrow |-5x + 2y + 19| = 18$$

$$\Rightarrow -5x + 2y + 19 = \pm 18$$

$$\Rightarrow -5x + 2y + 19 = 18 \text{ or } -5x + 2y + 19 = 18$$

$$\Rightarrow 5x - 2y - 1 = 0 \text{ or } 5x - 2y - 37 = 0$$

\therefore Locus of P is :

$$(5x - 2y - 1)(5x - 2y - 37) = 0.$$

8. **Find the equation of locus of a point which forms a triangle of area 2 with the point A(1, 1) and B(-2, 3).**

Sol. Same as above.

$$\text{Ans. } (2x + 3y - 1)(2x + 3y - 9) = 0$$

9. **If the distance from P to the points (2, 3) and (2, -3) are in the ratio 2 : 3, then find the equation of locus of P.**

Sol. Let P(x, y) be a point in locus.

Given points are A(2, 3), B(2, -3)

Given that $PA : PB = 2 : 3$

$$\Rightarrow 3PA = 2PB$$

$$\Rightarrow 9PA^2 = 4PB^2$$

$$\Rightarrow 9[(x-2)^2 + (y-3)^2] = 4[(x-2)^2 + (y+3)^2]$$

$$\Rightarrow 9[x^2 - 4x + 4 + y^2 - 6y + 9] = 4[x^2 - 4x + 4 + y^2 + 6y + 9]$$

$$\Rightarrow 5x^2 + 5y^2 - 20x - 78y + 65 = 0 \text{ which is the equation of locus.}$$

10. **A(1, 2), B(2, -3) and C(-2, 3) are three points. A point P moves such that $PA^2 + PB^2 = 2PC^2$. Show that the equation to the locus of P is $7x - 7y + 4 = 0$.**

Sol. Let P(x, y) be a point in locus.

Given points are A (1, 2), B (2, -3) and C(-2, 3)

Given that $PA^2 + PB^2 = 2PC^2$

$$\begin{aligned} \Rightarrow (x-1)^2 + (y-2)^2 + (x-2)^2 + (y+3)^2 &= 2[(x+2)^2 + (y-3)^2] \\ \Rightarrow 2x^2 + 2y^2 - 6x + 2y + 18 &= 2x^2 + 2y^2 + 8x - 12y + 26 \\ \Rightarrow 14x - 14y + 8 &= 0 \\ \Rightarrow 7x - 7y + 4 &= 0 \end{aligned}$$

Therefore, equation of locus is $7x - 7y + 4 = 0$

11. A straight rod of length 9 slides with its ends A, B always on the X and Y-axes respectively. Then find the locus of the centroid of ΔOAB .

Sol. The given rod AB meets X-axis at A and Y-axis at B.

Let $OA = a$ and $OB = b$ and $AB = 9$.

Coordinates of A are $(a, 0)$ and B are $(0, b)$.

Let $G(x, y)$ be the centroid of ΔOAB

But Coordinates of G of ΔOAB are $\left(\frac{a}{3}, \frac{b}{3}\right)$

Therefore, $\left(\frac{a}{3}, \frac{b}{3}\right) = (x, y)$


$$\Rightarrow \frac{a}{3} = x, \frac{b}{3} = y \Rightarrow a = 3x, b = 3y$$

But $OA^2 + OB^2 = AB^2$ and given $AB = 9$

$$\Rightarrow a^2 + b^2 = 81$$

$$\Rightarrow 9(x^2 + y^2) = 81$$

\therefore Equation to the locus of P is $x^2 + y^2 = 9$.



Problems for practice

1. Find the equation of the locus of a point which is at a distance 5 from $(-2, 3)$ in a plane.

Ans. $x^2 + y^2 + 4x - 6y - 12 = 0$.

2. Find the equation of locus of a point P, if the distance of P from $A(3, 0)$ is twice the distance of P from $B(-3, 0)$.

Ans. $x^2 + y^2 + 10x + 9 = 0$.

3. Find the locus of the third vertex of a right angled triangle, the ends of whose hypotenuse are $(4, 0)$ and $(0, 4)$.

Ans. $x^2 + y^2 - 4x - 4y = 0$

4. Find the equation of locus of P, if the ratio of the distances from P to $(5, -4)$ and $(7, 6)$ is $2 : 3$.

Ans. $5(x^2 + y^2) - 34x + 120y + 29 = 0$.

5. $A(2, 3)$ and $B(-3, 4)$ are two given points. Find the equation of locus of P so that the area of the triangle PAB is 8.5.

Ans. $x^2 + 10xy + 25y^2 - 34x - 170y = 0$