

## Random Variables and Probability Distributions

- \* If  $X : S \rightarrow R$  is a discrete random variable with range  $\{x_1, x_2, x_3, \dots\}$  then  $\sum_{r=1}^{\infty} P(X = x_r) = 1$
- \* Let  $X : S \rightarrow R$  be a discrete random variable with range  $\{x_1, x_2, x_3, \dots\}$ . If  $\sum x_r P(X = x_r)$  exists, then  $\sum x_r P(X = x_r)$  is called the mean of the random variable  $X$ . It is denoted by or  $\bar{x}$ .
- \* If  $\sum (x_r - \mu)^2 P(X = x_r)$  exists, then  $\sum (x_r - \mu)^2 P(X = x_r)$  is called variance of the random variable  $X$ . It is denoted by  $\sigma^2$ . The positive square root of the variance is called the standard deviation of the random variable  $X$ . It is denoted by  $\sigma$
- \* If the range of discrete random variable  $X$  is  $\{x_1, x_2, \dots, x_n, \dots\}$  and  $P(X = x_n) = P_n$  for every Integer  $n$  is given then  $\sigma^2 + \mu^2 = \sum x_n^2 P_n$

**Binomial Distribution:** A random variable  $X$  which takes values  $0, 1, 2, \dots, n$  is said to follow binomial distribution if its probability distribution function is given by

$$P(X = r) = {}^n C_r p^r q^{n-r}, r = 0, 1, 2, \dots, n \quad \text{where } p, q > 0 \text{ such that } p + q = 1.$$

- \* If the probability of happening of an event in one trial be  $p$ , then the probability of successive happening of that event in  $r$  trials is  $p^r$ .

Mean and variance of the binomial distribution

The mean of this distribution is  $\sum_{i=1}^n X_i p_i = \sum_{X=1}^n X \cdot {}^n C_X q^{n-X} p^X = np$ ,

The variance of the Binomial distribution is  $\sigma^2 = npq$  and the standard deviation is  $\sigma = \sqrt{npq}$ .

**The Poisson Distribution :** Let  $X$  be a discrete random variable which can take on the values  $0, 1, 2, \dots$  such that the probability function of  $X$  is given by

$$f(x) = P(X = x) = \frac{\lambda^x e^{-\lambda}}{x!}, \quad x = 0, 1, 2, \dots$$

where  $\lambda$  is a given positive constant. This distribution is called the Poisson distribution and a random variable having this distribution is said to be Poisson distributed.

### Very Short Answer Questions

1. A probability distribution function of a discrete random variable is zero except at the points  $x = 0, 1, 2$ . At these points it has the value

$p(0) = 3c^3, p(1) = 4c - 10c^2, p(2) = 5c - 1$  for some  $c > 0$ . Find the value of  $c$ .

Sol.

$$P(x = 0) + p(x = 1) + p(x = 2) = 1$$

$$3c^3 + 4c - 10c^2 + 5c - 1 = 1$$

$$3c^3 - 10c^2 + 9c - 2 = 0 \text{----- (1)}$$

$$\text{Put } c = 1, \text{ then } 3 - 10 + 9 - 2 = 12 - 12 = 0$$

$C = 1$  satisfy the above equation

$C = 1 \Rightarrow p(x = 1) = 4 - 10 = -6$  which is not possible.

Dividing (1) with  $c - 1$ ,

$$\text{We get } 3c^2 - 7c + 2 = 0$$

$$\Rightarrow (c - 2)(3c - 1) = 0$$

$$c = 2 \text{ or } c = 1/3$$

$c = 2 \Rightarrow p(x = 0) = 3 \cdot 2^3 = 24$  which is not possible

$$\therefore c = 1/3$$

2. Find the constant C, so that  $F(x) = C\left(\frac{2}{3}\right)^x, x = 1, 2, 3, \dots$  is the p.d.f of a discrete random variable X.

Sol. Given  $F(x) = C\left(\frac{2}{3}\right)^x, x = 1, 2, 3$

We know that  $p(x) = C\left(\frac{2}{3}\right)^x, x = 1, 2, 3 \dots$

$$\because \sum_{x=1}^{\infty} p(x) = 1 \Rightarrow \sum_{x=1}^{\infty} c\left(\frac{2}{3}\right)^x = 1$$

$$\Rightarrow c\left[\left(\frac{2}{3}\right) + \left(\frac{2}{3}\right)^2 + \left(\frac{2}{3}\right)^3 + \dots \infty\right] = 1$$

$$\Rightarrow C\frac{2}{3}\left[1 + \frac{2}{3} + \left(\frac{2}{3}\right)^2 + \dots \infty\right] = 1$$

$$\Rightarrow \frac{2c}{3}\left(\frac{1}{1 - \frac{2}{3}}\right) = 1 \Rightarrow \frac{2c}{3} \times 3 = 1 \Rightarrow c = \frac{1}{2}$$

3.

<b>X=x</b>	<b>-2</b>	<b>-1</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>
<b>P(X=x)</b>	<b>0.1</b>	<b>k</b>	<b>0.2</b>	<b>k</b>	<b>0.3</b>	<b>k</b>

Is the probability distribution of a random variable x. find the value of K and the variance of x.

Sol. We know that  $\sum_{i=1}^n p(x_i) = 1$

$$0.1 + k + 0.2 + k + 0.3 + k = 1$$

$$\Rightarrow 3k + 0.6 = 1$$

$$3k = 1 - 0.6 = 0.4 \Rightarrow k = \frac{0.4}{3} = 0.1333$$

$X=x$	-2	-1	0	1	2	3
$P(X=x)$	0.1	k	0.2	k	0.3	k
$X_i \cdot p(x_i)$	-0.2	-k	0	k	0.6	3k
$X_i^2 \cdot p(x_i)$	0.4	k	0	k	1.2	9k

$$\text{Mean} = \sum_{i=1}^n x_i p(x_i)$$

$$= -0.2 - k + 0 + 2k + 0.6 + 3k$$

$$= 4k + 0.4 = 4(0.1) + 0.4 = 0.4 + 0.4 = 0.8$$

$$\mu = 0.8$$

$$\text{Variance } (\sigma^2) = \sum_{i=1}^n x_i^2 p(x_i) - \mu^2$$

$$\therefore \text{Variance} = 4(0.1) + 1(k) + 0(0.2) + 1(2k) + 4(0.3) + 9k - \mu^2$$

$$= 0.4 + k + 0 + 2k + 4(0.3) + 9k - \mu^2$$

$$= 12k + 0.4 + 1.2 - (0.8)^2$$

$$= 12(0.1) + 1.6 - 0.64 = 1.2 + 1.6 - 0.64$$

$$\therefore \sigma^2 = 2.8 - 0.64 = 2.16$$

4.

$X=x$	-3	-2	-1	0	1	2	3
$P(X=x)$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$

Is the probability distribution of a random variable  $x$ . find the variance of  $x$ .

Sol.

$X=x$	-3	-2	-1	0	1	2	3
$P(X=x)$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{1}{9}$
$X_i \cdot p(x_i)$	$-\frac{3}{9}$	$-\frac{2}{9}$	$-\frac{1}{9}$	0	$\frac{1}{9}$	$\frac{2}{9}$	$\frac{3}{9}$
$X_i^2 \cdot p(x_i)$	$\frac{9}{9}$	$\frac{4}{9}$	$\frac{1}{9}$	0	$\frac{1}{9}$	$\frac{4}{9}$	$\frac{9}{9}$

$$\text{Mean } (\mu) = \sum_{i=1}^n x_i p(x_i)$$

$$= -\frac{3}{9} - \frac{2}{9} - \frac{1}{9} + 0 + \frac{1}{9} + \frac{2}{9} + \frac{3}{9} = (\mu) = 0$$

$$\text{Variance } (\sigma^2) = \sum_{i=1}^n x_i^2 p(x = x_i) - \mu^2$$

$$= \frac{9}{9} + \frac{4}{9} + \frac{1}{9} + 0 + \frac{1}{9} + \frac{4}{9} + \frac{9}{9} - 0^2 = \frac{28}{9} - 0$$

$$\sigma^2 = \frac{28}{9}$$

5. A random variable  $x$  has the following probability distribution.

$X=x$	0	1	2	3	4	5	6	7
$P(X=x)$	0	k	2k	2k	3k	$K^2$	$2k^2$	$7k^2+k$

Find i) k ii) the mean and iii)  $p(0 < x < 5)$ .

Sol.

We know that  $\sum_{i=1}^n p(x_i) = 1$

$$0 + k + 2k + 2k + 3k + K^2 + 2k^2 + 7k^2 + k = 1$$

$$\Rightarrow 10k^2 + 9k = 1 \Rightarrow 10k^2 + 9k - 1 = 0$$

$$\Rightarrow 10k^2 + 10k - k - 1 = 0$$

$$\Rightarrow 10k(k + 1) - 1(k + 1) = 0$$

$$\Rightarrow (10k - 1)(k + 1) = 0$$

$$K = \frac{1}{10}, -1 \text{ Since } k > 0 \therefore k = \frac{1}{10}$$

i)  $k = \frac{1}{10}$

ii)

$X=x$	0	1	2	3	4	5	6	7
$P(X=x)$	0	k	2k	2k	3k	$K^2$	$2k^2$	$7k^2+k$
$X_i.p(x_i)$	0	k	4k	6k	12k	$5k^2$	$12k^2$	$49k^2+7k$

$$\text{Mean} = \sum_{i=1}^n x_i p(x = x_i)$$

$$= 0 + k + 4k + 6k + 12k + 5k^2 + 12k^2 + 49k^2 + 7k$$

$$= 66k^2 + 30k$$

$$= 66\left(\frac{1}{100}\right) + 30 \times \left(\frac{1}{10}\right)$$

$$= 0.66 + 3 = 3.66$$

iii)  $p(0 < x < 5)$

$$p(0 < x < 5) =$$

$$p(x = 1) + p(x = 2) + p(x = 3) + p(x = 4)$$

$$= k + 2k + 2k + 3k = 8k$$

$$= 8 \frac{1}{10} = 8 \frac{1}{10} = \frac{4}{5}$$

6. In the experiment of tossing a coin  $n$  times, if the variable  $x$  denotes the number of heads and  $P(X = 4)$ ,  $P(X = 5)$ ,  $P(X = 6)$  are in arithmetic progression then find  $n$ .

Sol.  $X$  follows binomial distribution with

$$\text{Prob. Of getting head is } p = \frac{1}{2} \Rightarrow q = \frac{1}{2}$$

Given,  $P(X = 4)$ ,  $P(X = 5)$ ,  $P(X = 6)$  are in A.P

$$\Rightarrow {}^n C_4 \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^{n-4}, {}^n C_5 \left(\frac{1}{2}\right)^5 \left(\frac{1}{2}\right)^{n-5},$$

$${}^n C_6 \left(\frac{1}{2}\right)^6 \left(\frac{1}{2}\right)^{n-6} \text{ are in A.P}$$

$$\Rightarrow {}^n C_4, {}^n C_5, {}^n C_6 \text{ are in A.P}$$

$$\Rightarrow 2 {}^n C_5 = {}^n C_4 + {}^n C_6$$

$$\Rightarrow \frac{2(n!)}{5!(n-5)!} = \frac{n!}{4!(n-4)!} + \frac{n!}{6!(n-6)!}$$

$$\Rightarrow \frac{2(n!)}{5 \times 4!(n-5)(n-6)} = \frac{n!}{4!(n-4)(n-5)(n-6)!} + \frac{n!}{6 \times 5 \times 4!(n-6)!}$$

$$\Rightarrow \frac{2(n!)}{5 \times 4!(n-5)(n-6)!}$$

$$= \frac{n!}{4!(n-6)!} \left[ \frac{1}{(n-4)(n-5)} + \frac{1}{30} \right]$$

$$\Rightarrow \frac{2}{5(n-5)} = \frac{30 + (n-4)(n-5)}{30(n-4)(n-5)}$$

$$\Rightarrow 2 \times 30(n-4) = 5[30 + n^2 - 9n + 20]$$

$$\Rightarrow 12n - 48 = n^2 - 9n - 50$$

$$\Rightarrow n^2 - 21n + 98 = 0$$

$$\Rightarrow n^2 - 14n - 7n + 98 = 0$$

$$n(n-14) - 7(n-14) = 0$$

$$(n-7)(n-14) = 0$$

$$\therefore n = 7 \text{ or } 14$$

**7. Find the maximum number of times a fair coin must be tossed so that the probability of getting at least one head is at least 0.8.**

**Sol.** Let  $n$  be number of times a fair coin tossed  $x$  denotes the number of heads getting  $x$  follows binomial distribution with parameters  $n$  and  $p = 1/2$  given  $p(x \geq 1) \geq 0.8$

$$\Rightarrow 1 - p(x = 0) \geq 0.8 \Rightarrow p(x = 0) \leq 0.2$$

$$\Rightarrow {}^n C_0 \left(\frac{1}{2}\right)^n \leq 0.2 \Rightarrow \left(\frac{1}{2}\right)^n \leq \frac{1}{5}$$

The maximum value of  $n$  is 3.



8. The probability of a bomb hitting a bridge is  $1/2$  and three direct hits (not necessarily consecutive) are needed to destroy it. Find the minimum number of bombs required so that the probability of the bridge being destroyed is greater than 0.9.

**Sol.** Let  $n$  be the minimum number of bombs required and  $x$  be the number of bombs that hit the bridge, then  $x$  follows binomial distribution with parameters  $n$  and  $p = 1/2$ .

$$\text{Now } p(x \geq 3) > 0.9$$

$$\Rightarrow 1 - p(x < 3) > 0.9$$

$$\Rightarrow p(x < 3) < 0.1$$

$$\Rightarrow p(x = 0) + p(x = 1) + p(x = 2) < 0.1$$

$$\Rightarrow {}^n C_0 \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^n + {}^n C_1 \left(\frac{1}{2}\right)^1 \left(\frac{1}{2}\right)^{n-1} + {}^n C_2 \left(\frac{1}{2}\right)^2 \left(\frac{1}{2}\right)^{n-2} < 0.1$$

$$\Rightarrow 1 \cdot \frac{1}{2^n} + \frac{n}{2^n} + \frac{n(n-1)}{2} \frac{1}{2^2} < \frac{1}{10}$$

$$\Rightarrow 1 \cdot \frac{1}{2^n} + \frac{n}{2^n} + \frac{n^2 - n}{2 \cdot 2^n} < \frac{1}{10}$$

$$\Rightarrow \frac{1}{2^n} \left(1 + n + \frac{n^2 - n}{2}\right) < \frac{1}{10}$$

$$\Rightarrow \frac{1}{2^n} \left(\frac{2 + 2n + n^2 - n}{2}\right) < \frac{1}{10}$$

$$\Rightarrow 5(n^2 + n + 2) < 2^n$$

By trial and error, we get  $n \geq 9$

$\therefore$  The least value of  $n$  is 9

$\therefore n = 9$

9. If the difference between the mean and the variance of a binomial variate is  $5/9$  then, find the probability for the event of 2 successes, when the experiment is conducted 5 times.

Sol. Given  $n = 5$ , let  $p$  be the parameters of the binomial distribution

$$\text{Mean} - \text{Variance} = 5/9$$

$$np - npq = 5/9$$

$$np(1 - q) = 5/9, \quad \because p + q = 1$$

$$n.p^2 = 5/9 \Rightarrow 5p^2 = 5/9$$

$$p^2 = 1/9 \Rightarrow p = 1/3$$

$$q = 1 - p = 1 - 1/3 = 2/3$$

$$p(x = 2) = {}^5C_2 \left(\frac{2}{3}\right)^3 \left(\frac{1}{3}\right)^2 = 10 \times \frac{8}{27} \cdot \frac{1}{9} = \frac{80}{243}$$

$$\therefore \text{Prob. of the event of 2 success} = \frac{80}{243}$$

10. One in 9 ships is likely to be wrecked, when they are set on sail, when 6 ships are on sail, find the probability for (a) At least one will arrive safely (b) Exactly, 3 will arrive safely.

Sol.  $P$  = probability of ship to be wrecked =  $1/9$

$$q = 1 - p = 1 - \frac{1}{9} = \frac{8}{9}$$

Number of ships =  $n = 6$

$$p(x = 0) = {}^6C_0 \left(\frac{8}{9}\right)^{6-0} \left(\frac{1}{9}\right)^0 = \left(\frac{8}{9}\right)^6$$

a) Probability of at least one will arrive safely =  $p(x > 0) = 1 - p(x = 0)$

$$= 1 - \left(\frac{8}{9}\right)^6 = 1 - \frac{8^6}{9^6}$$

$$\begin{aligned} \text{b) } p(x = 3) &= {}^6C_3 \left(\frac{8}{9}\right)^3 \left(\frac{1}{9}\right)^3 \\ &= \frac{1}{9^6} \cdot {}^6C_3 \cdot 8^3 = 20 \left(\frac{8^3}{9^6}\right) \end{aligned}$$

**11. If the mean and variance of a binomial variable x are 2.4 and 1.44 respectively, find  $p(1 < x \leq 4)$ .**

**Sol.** Mean =  $np = 2.4$  ... (1)

Variance =  $npq = 1.44$  ... (2)

Dividing (2) by (1)

$$\frac{npq}{np} = \frac{1.44}{2.4}$$

$q = 0.6 = 3/5 \Rightarrow$

$p = 1 - q = 1 - 0.6 = 0.4 = 2/5$

Substituting in (1)

$n(0.4) = 2.4 \Rightarrow n = \frac{2.4}{0.4} = 6$

$P(1 < x \leq 4) = p(x = 2) + p(x = 3) + p(x = 4)$

$= {}^6C_2 q^4 \cdot p^2 + {}^6C_3 q^3 \cdot p^3 + {}^6C_4 q^2 \cdot p^4$

$= {}^6C_2 \left(\frac{3}{5}\right)^4 \left(\frac{2}{5}\right)^2 + {}^6C_3 \left(\frac{3}{5}\right)^3 \left(\frac{2}{5}\right)^3 + {}^6C_4 \left(\frac{3}{5}\right)^2 \left(\frac{2}{5}\right)^4 = \frac{6^2}{5^6} (15.9 + 20.6 + 15.4)$

$= \frac{36}{15625} (135 + 120 + 60) = \frac{315 \times 36}{15625}$

$= \frac{36 \times 63}{3125} = \frac{2268}{3125}$

12. It is given that 10% of the electric bulbs manufactured by a company are defective. In a sample of 20 bulbs, find the probability that more than 2 are defective.

Sol.  $p$  = probability of defective bulb =  $1/10$

$$q = 1 - p = 1 - \frac{1}{10} = \frac{9}{10}$$

$n$  = number of bulbs in the sample = 20

$$p(x > 2) = 1 - p(x \leq 2)$$

$$= 1 - [p(x = 0) + p(x = 1) + p(x = 2)]$$

$$p(x = 0) = {}^{20}C_0 \left(\frac{9}{10}\right)^{20} = \left(\frac{9}{10}\right)^{20}$$

$$p(x = 1) = {}^{20}C_1 \left(\frac{9}{10}\right)^{19} = \left(\frac{1}{10}\right) = \frac{20 \cdot 9^{19}}{10^{20}}$$

$$p(x = 2) = {}^{20}C_2 \left(\frac{9}{10}\right)^{18} = \left(\frac{1}{10}\right)^2 = \frac{190 \cdot 9^{18}}{10^{20}}$$

$$p(x > 2) = 1 - \left(\frac{9^{20}}{10^{20}} + \frac{20 \cdot 9^{19}}{10^{20}} + \frac{190 \cdot 9^{18}}{10^{20}}\right)$$

$$= 1 - \sum_{k=0}^2 {}^{20}C_k \left(\frac{9}{10}\right)^{20-k} \left(\frac{1}{10}\right)^k$$

$$= 1 - \sum_{k=0}^2 {}^{20}C_k \frac{9^{20-k}}{10^{20}} = \sum_{k=3}^{20} {}^{20}C_k \left(\frac{9^{20-k}}{10^{20}}\right)$$

13. On an average, rain falls on 12 days in every 30 days, find the probability that, rain will fall on just 3 days of a given week.

Sol. Given  $p = \frac{12}{30} = \frac{2}{5} \Rightarrow q = 1 - p = 1 - \frac{2}{5} = \frac{3}{5}$

$$n = 7, r = 3$$

$$p(x = 3) = {}^n C_r \cdot q^{n-r} \cdot p^r = {}^7 C_3 \left(\frac{3}{5}\right)^4 \left(\frac{2}{5}\right)^3$$

$$= 35 \cdot \left(\frac{3}{5}\right)^4 \left(\frac{2}{5}\right)^3 = \frac{35 \times 2^3 \times 3^4}{5^7}$$

14. For a binomial distribution with mean 6 and variance 2, find the first two terms of the distribution.

Sol. Let  $n, p$  be the parameters of a binomial distribution

$$\text{Mean } (np) = 6 \quad \dots(1)$$

$$\text{And variance } (npq) = 2 \quad \dots(2)$$

$$\text{then } \frac{npq}{np} = \frac{2}{6} \Rightarrow q = \frac{1}{3} \Rightarrow \because p = 1 - q = 1 - \frac{1}{3} = \frac{2}{3}$$

$$\text{From (1) } np = 6$$

$$n \left(\frac{2}{3}\right) = 6 \Rightarrow n = \frac{18}{2} = 9$$

First two terms of the distribution are

$$p(x = 0) = {}^9 C_0 \left(\frac{1}{3}\right)^9 = \frac{1}{3^9} \text{ and}$$

$$p(x = 1) = {}^9 C_1 \left(\frac{1}{3}\right)^8 \left(\frac{2}{3}\right) = \frac{2}{3^7}$$

15. In a city 10 accidents take place in a span of 50 days. Assuming that the number of accidents follows the poisson distribution, find the probability that there will be 3 or more accidents in a day.

Sol. Average number of accidents per day

$$\lambda = \frac{10}{50} = \frac{1}{5} = 0.2$$

The prob. That there will be 3 or more accidents in a day  $p(x \geq 3)$

$$\sum_{k=3}^{\infty} e^{-\lambda} \frac{\lambda^k}{k!}, \lambda = 0.2$$

### Short Answer Questions & Long Answer Questions

1. The range of a random variable  $x$  is  $\{0, 1, 2\}$ . Given that  $p(x = 0) = 3c^3$ ,  $p(x = 1) = 4c - 10c^2$ ,  $p(x = 2) = 5c - 1$

i) Find the value of  $c$

ii)  $p(x < 1)$ ,  $p(1 < x \leq 3)$

Sol.  $P(x = 0) + p(x = 1) + p(x = 2) = 1$

$$3c^3 + 4c - 10c^2 + 5c - 1 = 1$$

$$3c^3 - 10c^2 + 9c - 2 = 0$$

$C = 1$  satisfy this equation

$C = 1 \Rightarrow p(x = 0) = 3$  which is not possible dividing with  $c - 1$ , we get

$$3c^2 - 7c + 2 = 0 \Rightarrow (c - 2)(3c - 1) = 0$$

$d = 2$  or  $c = 1/3$

$c = 2 \Rightarrow p(x = 0) = 3 \cdot 2^3 = 24$  which is not possible

$\therefore c = 1/3$

**i)  $p(x < 1) = p(x = 0)$**

$$= 3.c^3 = \left(\frac{1}{3}\right)^3 = 3 \cdot \frac{1}{27} = \frac{1}{9}$$

**ii)  $p(1 < x \leq 2) = p(x = 2) = 5c - 1$**

$$= \frac{5}{3} - 1 = \frac{2}{3}$$

**iii)  $p(0 < x \leq 3) = p(x = 1) + p(x = 2)$**

$$= 4c - 10c^2 + 5c - 1$$

$$= 9c - 10c^2 - 1 = 9 \cdot \frac{1}{3} - 10 \cdot \frac{1}{9} - 1$$

$$= 3 - \frac{10}{9} - 1 = 2 - \frac{10}{9} = \frac{8}{9}$$

**2. The rang of a random variable x is {1, 2, 3...} and  $p(x = k) = \frac{c^k}{k!}$**

**(k = 1, 2, 3, .....)** Find the value of C and  $p(0 < x < 3)$

**Sol.** Sum of the probabilities = 1

$$\sum p(x = k) = 1 \Rightarrow \sum_{k=1}^{\infty} \frac{c^k}{k!} = 1$$

$$\Rightarrow c + \frac{c^2}{2} + \frac{c^3}{3} + \dots = 1$$

Adding 1 on both sides

$$\Rightarrow 1 + c + \frac{c^2}{2} + \frac{c^3}{3} + \dots = 2$$

$$\Rightarrow e^c = 2 \Rightarrow \log_e e^c = \log_e 2$$

$$\Rightarrow c = \log_e 2$$

$$P(0 < x < 3) = p(x = 1) + p(x = 2)$$

$$= c + \frac{c^2}{2} = \log_e 2 + \frac{(\log_e 2)^2}{2}$$

**3. Five coins are tossed 320 times. Find the frequencies of the distribution of number of heads and tabulate the result.**

**Sol.** 5 coins are tossed 320 times

Prob. of getting a head on a coin

$$p = \frac{1}{2}, n = 5$$

Prob. of having x heads

$$p(x = x) = {}^5C_x (q)^5 \left(\frac{1}{2}\right)^x$$
$$= {}^5C_x \left(\frac{1}{2}\right)^{5-x} \left(\frac{1}{2}\right)^x = {}^5C_x \left(\frac{1}{2}\right)^5 \quad x = 0, 1, 2, 3, 4, 5$$

Frequencies of the distribution of number of heads = N.P(X = x)

$$= 320 \left[ {}^5C_x \left(\frac{1}{2}\right)^5 \right]; x = 0, 1, 2, 3, 4, 5$$

Frequency of

$$\text{Having 0 head} = 320 \times {}^5C_0 \times \left(\frac{1}{2}\right)^5 = 10$$

$$\text{Having 1 head} = 320 \times {}^5C_1 \times \left(\frac{1}{2}\right)^5 = 50$$



$$\text{Having 2 head} = 320 \times {}^5C_2 \times \left(\frac{1}{2}\right)^5 = 100$$

$$\text{Having 3 head} = 320 \times {}^5C_3 \times \left(\frac{1}{2}\right)^5 = 100$$

$$\text{Having 4 head} = 320 \times {}^5C_4 \times \left(\frac{1}{2}\right)^5 = 50$$

$$\text{Having 5 head} = 320 \times {}^5C_5 \times \left(\frac{1}{2}\right)^5 = 10$$

<b>N(H)</b>	0	1	2	3	4	5
<b>f</b>	10	50	100	100	50	10

**4. Find the probability of guessing at least 6 out of 10 of answers in**

**(i) True or false type examination    ii) Multiple choice with 4 possible answers.**

**Sol.i)** Since the answers are in true or false type.

$$\text{Prob. of success } p = \frac{1}{2}, q = \frac{1}{2}$$

Prob. of guessing at least 6 out of 10

$$p(x \geq 6) = \sum_6^{10} {}^{10}C_k \left(\frac{1}{2}\right)^{10-k} \left(\frac{1}{2}\right)^k = \sum_6^{10} {}^{10}C_k \left(\frac{1}{2}\right)^{10}$$

ii) Since the answers are in multiple choice with 4 possible answers

$$\text{Prob. of success } p = 1/4, q = 3/4$$

Prob. of guessing at least 6 out of 10

$$p(x \geq 6) = \sum_6^{10} {}^{10}C_k \left(\frac{1}{4}\right)^k \left(\frac{3}{4}\right)^{10-k}$$

5. The number of persons joining a cinema ticket counter in a minute has poisson distribution with parameter 6. Find the probability that i) no one joins the queue in a particular minute ii) two or more persons join the queue in a minute.

Sol. Here  $\lambda = 6$

i) prob. That no one joins the queue in a particular minute

$$p(x = 0) = \frac{e^{-\lambda} \lambda^0}{0!} = e^{-6}$$

ii) prob. that two or more persons join the queue in a minute

$$\begin{aligned} p(x \geq 2) &= 1 - p(x \leq 1) \\ &= 1 - [p(x = 0) + p(x = 1)] \\ &= 1 - \left[ e^{-\lambda} \frac{\lambda^0}{0!} + e^{-\lambda} \frac{\lambda^1}{1!} \right] \\ &= 1 - \left[ e^{-6} + \frac{e^{-6}(6)}{1!} \right] = 1 - 7e^{-6} \end{aligned}$$

6. A cubical die is thrown. Find the mean and variance of x, giving the number on the face that shows up.

Sol. Let S be the sample space and x be the random variable associated with S, where p(x) is given by the following table

$X=x_i$	1	2	3	4	5	6
$P(X=x_i)$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$
$X_i.p(x_i)$	$1/6$	$2/6$	$3/6$	$4/6$	$5/6$	$6/6$
$X_i^2.p(x_i)$	$1/6$	$4/6$	$9/6$	$16/6$	$25/6$	$36/6$

Mean of  $x = \mu = \sum(X = x_i) P(X = x_i)$

$$= 1\left(\frac{1}{6}\right) + 2\left(\frac{1}{6}\right) + 3\left(\frac{1}{6}\right) + 4\left(\frac{1}{6}\right) + 5\left(\frac{1}{6}\right) + 6\left(\frac{1}{6}\right)$$

$$= \frac{1}{6}(1+2+3+4+5+6)$$

$$= \frac{1}{6} \frac{(6)(6+1)}{2} = \frac{7}{2} = 3.5$$

Variance of  $x = \sigma^2 = \sum x_i^2 p(X = x_i) - \mu^2$

$$= \frac{1}{6}(1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2) - \frac{49}{4}$$

$$= \frac{1}{6} \frac{(6)(6+1)(2 \times 6 + 1)}{6} - \frac{49}{4}$$

$$= \frac{91}{6} - \frac{49}{4} = \frac{182 - 147}{12} = \frac{35}{12}$$

7. The probability distribution of a random variable  $x$  is given below. Find the value of  $k$ , and the mean and variance of  $x$

$X=x_i$	1	2	3	4	5
$P(X=x_i)$	$k$	$2k$	$3k$	$4k$	$5k$

Sol. we have  $\sum_{r=1}^5 p(X = x_r) = 1$

$$\Rightarrow k + 2k + 3k + 4k + 5k = 1 \Rightarrow k = \frac{1}{15}$$

$X=x_i$	1	2	3	4	5
$P(X=x_i)$	$k$	$2k$	$3k$	$4k$	$5k$
$X_i \cdot p(x_i)$	$k$	$4k$	$9k$	$16k$	$25k$
$X_i^2 \cdot p(x_i)$	$k$	$8k$	$27k$	$64k$	$125k$

$$\begin{aligned} \text{Mean } \mu \text{ of } x &= \sum_{r=1}^5 r p(x = x_r) = \sum_{r=1}^5 r(rk) \\ &= 1.(k) + 2.(2k) + 3.(3k) + 4.(4k) + 5.(5k) \\ &= 55k \\ &= 55 \times \frac{1}{15} = \frac{11}{3} \end{aligned}$$

$$\begin{aligned} \text{Variance } (\sigma^2) &= \sum_{i=1}^n x_i^2 p(x = x_i) - \mu^2 \\ &= k + 8k + 27k + 64k + 125k - \left(\frac{11}{3}\right)^2 \\ &= 225k - \frac{121}{9} = 225 \times \frac{1}{15} - \frac{121}{9} \\ &= \frac{135 - 121}{9} = \frac{14}{9} \end{aligned}$$

8. If  $X$  is a random variable with the probability distribution.  $P(X = k) = \frac{(k+1)c}{2^k}$ , ( $k = 0, 1, 2, 3, \dots$ ) then find  $C$ .

Sol. given  $p(x = k) = \frac{(k+1)c}{2^k}$  ( $k = 0, 1, 2, 3, \dots$ )

$$\sum_{k=0}^{\infty} p(x = k) = 1$$

$$\sum_{k=0}^{\infty} \frac{(k+1)c}{2^k} = c \left( 1 + 2 \cdot \frac{1}{2} + 3 \left(\frac{1}{2}\right)^2 + k\alpha \right) = 1$$

$$c \left[ \frac{1}{1 - \frac{1}{2}} + \frac{1 \cdot \frac{1}{2}}{\left(1 - \frac{1}{2}\right)^2} \right] = 1$$

**Hint:** in A.G.P.  $S_{\infty} = \frac{a}{1-r} + \frac{dr}{(1-r)^2}$

Here  $a = 1$ ,  $d = 1$ ,  $r = 1/2$

$$c \left[ \frac{1}{\frac{1}{2}} + \frac{\frac{1}{2}}{\left(\frac{1}{2}\right)^2} \right] = 1$$

$$c [2 + 2] = 1$$

$$\therefore c = \frac{1}{4}$$

**9. Let  $x$  be a random variable such that  $p(x = -2) = p(x = -1) = p(x = 2) = p(x = 3) = 1/6$  and  $p(x = 0) = 1/3$ . Find the mean and variance of  $x$ .**

**Sol.** Mean

$$= (-2) \frac{1}{6} + (-1) \frac{1}{6} + 2 \left( \frac{1}{6} \right) + (1) \left( \frac{1}{6} \right) + 0 \cdot \left( \frac{1}{3} \right)$$

$$= -\frac{2}{6} - \frac{1}{6} + \frac{2}{6} + \frac{1}{6} + 0$$

$$\mu = 0$$

$$\text{Variance } (\sigma^2) = (-2)^2 \left( \frac{1}{6} \right) + (-1)^2 \left( \frac{1}{6} \right)$$

$$- 0^2 \left( \frac{1}{3} \right) + 2^2 \left( \frac{1}{6} \right) + 1^2 \left( \frac{1}{6} \right)$$

$$= \frac{4}{6} + \frac{1}{6} + 0 + \frac{4}{6} + \frac{1}{6}$$

$$= \frac{10}{6} - \frac{5}{3}$$

**10. Two dice are rolled at random. Find the probability distribution of the sum of the numbers on them. Find the mean of the random variable.**

**Sol.** When two dice are rolled, the sample space  $S$  contains  $6 \times 6 = 36$  sample points.

$$S = \{(1, 1), (1, 2), \dots, (1, 6), (2, 1), (2, 2), \dots, (6, 6)\}$$

Let  $x$  denote the sum of the numbers on the two dice

Then the range  $x = \{2, 3, 4, \dots, 12\}$

Probability Distribution of  $x$  is given by the following table.

$X=x_i$	2	3	4	5	6	7	8	9	10	11	12
$P(X=x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$
$X_i \cdot p(x_i)$	$2/36$	$6/36$	$12/36$	$20/36$	$30/36$	$42/36$	$40/36$	$36/36$	$30/36$	$22/36$	$12/36$

$$\text{Mean of } x = \mu = \sum_{i=1}^{12} x_i p(X = x_i)$$

$$= 2 \cdot \frac{1}{36} + 3 \cdot \frac{2}{36} + 4 \cdot \frac{3}{36} + 5 \cdot \frac{4}{36} + 6 \cdot \frac{5}{36} + 7 \cdot \frac{6}{36} + 8 \cdot \frac{5}{36} + 9 \cdot \frac{4}{36} + 10 \cdot \frac{3}{36} + 11 \cdot \frac{2}{36} + 12 \cdot \frac{1}{36}$$

$$= \frac{1}{36} (2 + 6 + 12 + 20 + 30 + 42 + 40 + 36 + 30 + 22 + 12)$$

$$= \frac{252}{36} = 7$$

**11. 8 coins are tossed simultaneously. Find the probability of getting at least 6 heads.**

**Sol.**  $p$  = probability of getting head =  $\frac{1}{2}$

$$q = 1 - p = 1 - \frac{1}{2} = \frac{1}{2}; n = 8$$

$$p(x \geq 6) = p(x = 6) + p(x = 7) + p(x = 8)$$

$$\begin{aligned}
 &= {}^8C_6 \left(\frac{1}{2}\right)^6 \left(\frac{1}{2}\right)^2 + {}^8C_7 \left(\frac{1}{2}\right)^7 \left(\frac{1}{2}\right)^1 + {}^8C_8 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^0 \\
 &= \left(\frac{1}{2}\right)^8 [{}^8C_6 + {}^8C_7 + {}^8C_8] \\
 &= \frac{1}{256} [28 + 8 + 1] = \frac{37}{256}
 \end{aligned}$$

**12. The mean and variance of a binomial distribution are 4 and 3 respectively. Fix the distribution and find  $p(x \geq 1)$**

**Sol.** Given distribution is binomial distribution with mean =  $np = 4$

$$\text{Variance} = npq = 3$$

$$\therefore \frac{npq}{np} = \frac{3}{4} \Rightarrow q = \frac{3}{4}$$

$$\text{So that } p = 1 - q = 1 - \frac{3}{4} = \frac{1}{4}$$

$$\therefore np = 4$$

$$n \frac{1}{4} = 4 \Rightarrow n = 16$$

$$P(x \geq 1) = 1 - p(x = 0)$$

$$= 1 - {}^{16}C_0 \left(\frac{1}{4}\right)^0 \left(\frac{3}{4}\right)^{16-0} = 1 - \left(\frac{3}{4}\right)^{16}$$

$$\therefore p(x \geq 1)$$

$$= 1 - \left(\frac{3}{4}\right)^{16}$$

13. The probability that a person chosen at Random is left handed (in hand writing) is 0.1. What is the probability that in a group of ten people there is one, who is left handed?

Sol. Here  $n = 10$

$$P = 0.1$$

$$q = 1 - p = 1 - 0.1 = 0.9$$

$$\begin{aligned} p(x = 1) &= {}^{10}C_1 (0.1)^1 (0.9)^{10-1} \\ &= 10 \times 0.1 \times (0.9)^9 \\ &= 1 \times (0.9)^9 = (0.9)^9 \end{aligned}$$

14. In a book of 450 pages, there are 400 typographical errors. Assuming that following the poisson law, the number of errors per page, find the probability that a random sample of 5 pages will contain no typographical error?

Sol. The average number of errors per page in the book is

$$\lambda = \frac{400}{450} = \frac{8}{9}$$

Here  $r = 0$

$$p(x = r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

$$p(x = 0) = \frac{e^{-8/9} \left(\frac{8}{9}\right)^0}{0!} = e^{-8/9}$$

∴ The required probability that a random sample of 5 pages will contain no error is

$$[p(x = 0)]^5 = \left(e^{-8/9}\right)^5$$



15. Deficiency of red cells in the blood cells is determined by examining a specimen of blood under a microscope. Suppose a small fixed volume contains on an average 20 red cells for normal persons. Using the poisson distribution, find the probability that a specimen of blood taken from a normal person will contain less than 15 red cells.

Sol. Here  $\lambda = 20$

$$p(x < 15) = \sum_{r=0}^{14} p(x = r)$$
$$= \sum_{r=0}^{14} \frac{e^{-\lambda} \lambda^r}{r!} = \sum_{r=0}^{14} e^{-20} \frac{20^r}{r!}$$

16. A poisson variable satisfies  $p(x = 1) = p(x = 2)$ . Find  $p(x = 5)$

Sol. Given  $p(x = 1) = p(x = 2)$

$$p(x = r) = \frac{\lambda^r e^{-\lambda}}{r!}, \lambda > 0$$

$$\frac{\lambda^1 e^{-\lambda}}{1!} = \frac{\lambda^2 e^{-\lambda}}{2!}$$

$$\lambda = 2, (\because \lambda > 0)$$

$$\therefore p(x = 5) = \frac{2^5 e^{-2}}{5!}$$

$$= \frac{32}{120e^2} = \frac{4}{15e^2}$$