

INVERSE TRIGONOMETRIC FUNCTIONS

PREVIOUS EAMCET BITS

1. $\cos^{-1}\left(\frac{-1}{2}\right) - 2\sin^{-1}\left(\frac{1}{2}\right) + 3\cos^{-1}\left(\frac{-1}{\sqrt{2}}\right) - 4\tan^{-1}(-1) =$ [EAMCET 2009]

- 1) $\frac{19\pi}{12}$ 2) $\frac{35\pi}{12}$ 3) $\frac{47\pi}{12}$ 4) $\frac{43\pi}{12}$

Ans: 4

Sol. $\frac{2\pi}{3} - 2 \times \frac{\pi}{6} + 3 \times \frac{3\pi}{4} + 4 \times \frac{\pi}{4} = \frac{43\pi}{12}$

2. If $\sin^{-1}\left(\frac{3}{x}\right) + \sin^{-1}\left(\frac{4}{x}\right) = \frac{\pi}{2}$ then $x =$ [EAMCET 2008]

- 1) 3 2) 5 3) 7 4) 11

Ans: 2

Sol. $\sin^{-1}\left(\frac{3}{x}\right) + \sin^{-1}\left(\frac{4}{x}\right) = \frac{\pi}{2} \Rightarrow \sin^{-1}\frac{3}{x} = \cos^{-1}\frac{4}{x}$
 $\Rightarrow \sin^{-1}\frac{3}{x} = \sin^{-1}\sqrt{1 - \frac{16}{x^2}} \Rightarrow \frac{9}{x^2} = 1 - \frac{16}{x^2} \Rightarrow x^2 = 25$

3. The value of x where $x > 0$ and $\tan\left(\sec^{-1}\left(\frac{1}{x}\right)\right) = \sin\left(\tan^{-1} 2\right)$ is [EAMCET 2007]

- 1) $\sqrt{5}$ 2) $\frac{\sqrt{5}}{3}$ 3) 1 4) $\frac{2}{3}$

Ans: 2

Sol. $\tan\left(\sec^{-1}\frac{1}{x}\right) = \sin\left(\tan^{-1} 2\right)$
 $\tan\left(\tan^{-1}\frac{\sqrt{1-x^2}}{x}\right) = \sin\left(\sin^{-1}\frac{2}{\sqrt{5}}\right)$
 $\Rightarrow \frac{\sqrt{1-x^2}}{x} = \frac{2}{\sqrt{5}} \Rightarrow x = \frac{\sqrt{5}}{3}$

4. $\sin^{-1}\frac{4}{5} + 2\tan^{-1}\frac{1}{3} =$ [EAMCET 2005]

- 1) $\frac{\pi}{3}$ 2) $\frac{\pi}{4}$ 3) $\frac{\pi}{2}$ 4) 0

Ans: 3

Sol. $\sin^{-1}\frac{4}{5} + 2\tan^{-1}\frac{1}{3} = \sin^{-1}\frac{4}{5} + \tan^{-1}\frac{3}{4}$
 $= \sin^{-1}\frac{4}{5} + \cos^{-1}\frac{4}{5} = \frac{\pi}{2}$

5. $\sin^{-1}x + \sin^{-1}(1-x) = \cos^{-1}x \Rightarrow x \in$ [EAMCET 2004]

1) $\{1, 0\}$

2) $\{-1, 1\}$

3) $\left\{0, \frac{1}{2}\right\}$

4) $\{2, 0\}$

Ans: 3

Sol. $\sin^{-1}(1-x) = \frac{\pi}{2} - 2\sin^{-1}(x)$

$$1-x = 1-2x^2 \Rightarrow x = 0, \frac{1}{2}$$

6. $\cos\left[\cos^{-1}\left(\frac{1}{7}\right) + \sin^{-1}\left(-\frac{1}{7}\right)\right] = \dots\dots$

[EAMCET 2003]

1) $-\frac{1}{3}$

2) 0

3) $\frac{1}{3}$

4) $\frac{4}{9}$

Ans: 2

Sol. $\cos\left[\cos^{-1}\left(-\frac{1}{7}\right) + \sin^{-1}\left(-\frac{1}{7}\right)\right]$

$$= \cos\frac{\pi}{2} = 0 \quad [\because \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}]$$

7. If $\sin^{-1} x - \cos^{-1} x = \frac{\pi}{6}$, then $x =$

1) $\frac{1}{2}$

2) $\frac{\sqrt{3}}{2}$

3) $-\frac{1}{2}$

4) $-\frac{\sqrt{3}}{2}$

Ans: 2

Sol. By verification $x = \sqrt{3}/2$

8. $\sec^2(\tan^{-1} 2) + \csc^2(\cot^{-1} 3) =$

1) 5

2) 10

3) 15

[EAMCET 2001]

Ans: 3

Sol. Let $\tan^{-1}(2) = \alpha$ and $\cot^{-1}(3) = \beta$

$\tan \alpha = 2; \cot \beta = 3$

$$\Rightarrow \sec \alpha = \sqrt{5}; \csc \beta = \sqrt{10}$$

$$\therefore \sec^2 \alpha + \csc^2 \beta = 5 + 10 = 15$$

9. If $\tan^{-1} 3 + \tan^{-1} x = \tan^{-1} 8$, then $x =$

1) 5

2) 1/5

3) 5/14

[EAMCET 2000]

Ans: 2

Sol. $\tan^{-1} 3 + \tan^{-1} x = \tan^{-1} 8$

$$\frac{3+x}{1-3x} = 8 \Rightarrow x = \frac{1}{5}$$

4) 14/5

விடைகள்