

# TANGENT AND NORMALS

## PREVIOUS EAMCET BITS

1. The equation to the normal to the curve  $y^4 = ax^3$  at  $(a, a)$  is **[EAMCET 2008]**  
 1)  $x + 2y = 3a$       2)  $3x - 4y + a = 0$       3)  $4x + 3y = 7a$       4)  $4x - 3y = 0$

Ans: 3

Sol.  $y^4 = ax^3 \Rightarrow 4y^3 \frac{dy}{dx} = 3ax^2 \Rightarrow \frac{3ax^2}{4y^3} \Rightarrow m = \left( \frac{dy}{dx} \right)_{(a,a)} = \frac{3a^3}{4a^3} = \frac{3}{4}$

Equation of the normal is  $y - a = -\frac{4}{3}(x - a) \Rightarrow 4x + 3y = 7a$

2. The length of the subtangent at  $(2, 2)$  to the curve  $x^5 = 2y^4$  is **[EAMCET 2008]**  
 1)  $5/2$       2)  $8/5$       3)  $2/5$       4)  $5/8$

Ans: 2

Sol.  $x^5 = 2y^4 \Rightarrow 5x^4 = 8y^3 \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{5x^4}{8y^3} = \frac{5(2)^4}{8(2)^3} = \frac{5}{4}$

Length of the subtangent  $\frac{2}{m} = \frac{2}{5/4} = \frac{8}{5}$

3. The length of tangent, subtangent normal and subnormal for the curve  $y = x^2 + x - 1$  at  $(1, 1)$  are A, B, C and D respectively, then their increasing order is **[EAMCET 2007]**  
 1) B, D, A, C      2) B, A, C, D      3) A, B, C, D      4) B, A, D, C

Ans: 4

Sol.  $m = \left( \frac{dy}{dx} \right)_{(1,1)} = 3$

A = length of tangent =  $\frac{\sqrt{10}}{3}$

B = length of subtangent =  $\frac{1}{3}$

C = length of normal =  $\sqrt{10}$

D = length of subnormal = 3

$\therefore B < A < D < C$

4. If  $\theta$  is the angle between the curves  $xy = 2$  and  $x^2 + 4y = 0$ , then  $\tan\theta =$  **[EAMCET 2006]**  
 1) 1      2) -1      3) 2      4) 3

Ans: 4

Sol.  $y = \frac{2}{x}, \quad y = \frac{-x^2}{4}$

$x^3 = -8 \quad (-2, -1)$

$x = -2$

$y = -1$

$m_1 = \frac{dy}{dx} = \frac{-2}{x^2} = \frac{-1}{2}$

$$m_2 = \frac{dy}{dx} = \frac{-2x}{4} = \frac{-x}{2} = 1$$

$$\tan \theta = \left| \frac{m_1 + m_2}{1 + m_1 m_2} \right| = \left| \frac{\frac{-1}{2} - 1}{1 - \frac{1}{2}} \right| = \left| \frac{\frac{-3}{2}}{\frac{1}{2}} \right| = 3$$

5. Match the points on the curve  $2y^2 = x + 1$  with the slopes of normals at those points and choose the correct answer. **[EAMCET 2004]**

List - I

I) (7, 2)

II)  $\left(0, \frac{1}{\sqrt{2}}\right)$ 

III) (1, -1)

IV)  $(3, \sqrt{2})$ 

List - II

a)  $-4\sqrt{2}$ 

b) -8

c) 4

d) 0

e)  $-2\sqrt{2}$ 

1) b, d, c, a

2) b, e, c, a

3) b, c, e, a

4) b, e, a, c

Ans: 2

Sol.  $2y^2 = x + 1 \Rightarrow \frac{-dx}{dy} = -4y = \text{slope of the normal}$

$$-\frac{dx}{dy} \text{ at } (7, 2) = -8$$

$$-\frac{dx}{dy} \text{ at } \left(0, \frac{1}{\sqrt{2}}\right) = -2\sqrt{2}$$

$$-\frac{dx}{dy} \text{ at } (1, -1) = 4;$$

$$-\frac{dx}{dy} \text{ at } (3, \sqrt{2}) = -4\sqrt{2}$$

6. The angle between the curves  $y = \sin x$  and  $y = \cos x$  is

1)  $\tan^{-1}(2\sqrt{2})$ 2)  $\tan^{-1}(3\sqrt{2})$ 3)  $\tan^{-1}(3\sqrt{3})$ 4)  $\tan^{-1}(5\sqrt{2})$ **[EAMCET 2003]**

Ans: 1

Sol.  $y = \sin x; y = \cos x \Rightarrow x = \frac{\pi}{4}$

$$m_1 = \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}$$

$$m_2 = -\sin \frac{\pi}{4} = \frac{-1}{\sqrt{2}}$$

$$\theta = \tan^{-1} \left( \frac{m_1 - m_2}{1 + m_1 m_2} \right) = \tan^{-1} (2\sqrt{2})$$

7. The two curves  $x = y^2$ ,  $xy = a^3$  cut orthogonally at a point, then  $a^2 =$  [EAMCET 2002]  
 1)  $1/3$                       2)  $1/2$                       3)  $2$                       4)  $3$

Ans: 2

Sol.  $x = y^2$ ,  $xy = a^3$

$$2ym_1 = 1 \quad m_2 = \frac{-a^3}{x^2} \Rightarrow m_1 = \frac{1}{2y}$$

$$m_1 m_2 = -1 \Rightarrow \frac{1}{2y} \cdot \frac{a^3}{x^2} = +1$$

$$\frac{xy}{2x^2y} = 1 \Rightarrow x = \frac{1}{2}$$

$$a^6 = x^2 y^2 = x^3 = \frac{1}{8} \Rightarrow (a^2)^3 = \left(\frac{1}{2}\right)^3 \therefore a^2 = \frac{1}{2}$$

8. The equation of the tangent to the curve  $6y = 7 - x^3$  at  $(1, 1)$  is [EAMCET 2001]  
 1)  $2x + y = 3$               2)  $x + 2y = 3$               3)  $x + y = -1$               4)  $x + y + 2 = 0$

Ans: 2

Sol.  $6y = 7 - x^3 \Rightarrow \frac{dy}{dx}$  at  $(1,1) = -\frac{1}{2}$

$$\text{Equation of the tangent is } y - 1 = \frac{-1}{2}(x - 1)$$

$$\Rightarrow x + 2y - 3 = 0$$

9. The angle between the curves  $y^2 = 4x$ ,  $x^2 = 4y$  at  $(4, 4)$  is [EAMCET 2000]  
 1)  $\tan^{-1}\left(\frac{1}{2}\right)$               2)  $\tan^{-1}\left(\frac{3}{4}\right)$               3)  $\frac{\pi}{2}$                       4)  $\frac{\pi}{4}$

Ans: 2

Sol.  $y^2 = 4x \Rightarrow \frac{dy}{dx}$  at  $(4,4) = \frac{1}{2} = m_1$

$$x^2 = 4y \Rightarrow \frac{dy}{dx}$$
 at  $(4,4) = 2 = m_2$

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| \Rightarrow \theta = \tan^{-1}\left(\frac{3}{4}\right)$$

10. Area of the triangle formed by the normal to the curve  $x = e^{\sin y}$  at  $(1, 0)$  with the coordinate axes is [EAMCET 2000]

- 1)  $\frac{1}{4}$                       2)  $\frac{1}{2}$                       3)  $\frac{3}{4}$                       4)  $1$

Ans: 2

Sol.  $x = e^{\sin y} \Rightarrow \log x = \sin y \Rightarrow \frac{dy}{dx} = \frac{1}{x \cos y}$

$$\frac{dy}{dx}$$
 at  $(1,0) = 1 = \text{slope of the tangent}$

$$\therefore \text{Equation of the normal is } x + y - 1 = 0$$

$$\therefore \text{Area of the triangle with normal and coordinate axes} = 1/2$$

\* \* \*