

Alkali Metals

Synopsis

- * Lithium (Li), Sodium (Na), Potassium (K), Rubidium (Rb), Cesium (Cs) and Francium (Fr) are called Alkali metals.
- * Francium is highly radioactive. Its longest-lived isotope ^{223}Fr has a half-life of only 21 minutes.
- * Oxides of these metals dissolve in water giving strong alkalis. So these elements are known as alkali metals.
- * Abundance: The abundance in earth crust decreases with the increase in atomic weight.
- * Electronic configuration: These elements belong to s-block and their valence shell configuration is ns^1 .
- * No element contains 18 electrons in their (n-1) shell.
- * Atomic volume: Atomic volume of alkali metals is highest in each period and goes on increasing down the group

Element	Li	Na	K	Rb	Cs
GMV in cm^3	13	24	46	56	71

- * Density: These elements have low densities when compared with the other metals.
- * Potassium is less dense than sodium due to
 1. Sudden large increase in the atomic size from Na to K.
 2. The presence of vacant d-orbitals
 3. Large interatomic distances in the crystal lattice.
- * Most electro positive element is Cs.
- * Hardness: These are soft metals. Softness further increases down the group due to decrease in the strength of metallic bond.
- * Valency & oxidation states: The alkali metal atoms show only +1 oxidation state, because their unipositive ions have the stable inert gas configuration ($s^2 \text{ or } s^2 P^6$) in the valence shell.
- * **Nature:** Alkali metal ions (in the form of salts) are colorless and diamagnetic, since all the electrons are paired. Some of their compounds like $\text{K}_2\text{Cr}_2\text{O}_4$, KMnO_4 are colored which is due to their oxyanions.

- * Second ionization potential values of alkali metals are very high, because the second electron is to be removed from the ion with stable inert gas configuration.
- * **Flame Test:** Alkali metals and their salts give flame colouration when heated with conc. HCl in Bunsen flame.
- * Exhibit colour due to absorption of visible light.
- * The characteristic flame colours of alkali metals are

Metal	Colour	λ in \AA
Li	Crimson red	6708
Na	Golden Yellow	5890
K	Lilac Blue	4404
Rb	Red-Violet	4202
Cs	Blue-Violet	4556


- * The melting points order is
 $Li > Na > K > Rb > Cs$
- * The boiling points order is
 $Li > Na > K > Cs > Rb$
- * The increasing order of SRP values of alkali metals is $Li < Cs < Rb < K < Na$
(SRP values -ve)
- * The hydration enthalpies of alkali metal ions decrease with increase in ionic sizes.
 $Li^+ > Na^+ > K^+ > Rb^+ > Cs^+$
- * Alkali metals when dissolved in liquid ammonia in the absence of impurities form a deep blue colored solution.
$$M + (x + y)NH_3 \rightarrow [M(NH_3)_x]^+ + [e(NH_3)_y]^-$$
- * The blue colour of the solution is due to ammoniated (solvate) electrons.
(**Note:** In concentrated solution, the blue colour changes to bronze colour and become diamagnetic due to cluster formation).
- * The solutions are paramagnetic and on standing slowly liberated hydrogen resulting in the formation of amide.
$$M_{(am)}^+ + e^- + NH_3(liq) \rightarrow MNH_{2(am)} + 1/2 H_2(g)$$
- * The electrical conductivity of these solutions is by the movement of ions as well as electrons. Hence they are called mixed conductors.

- * Due to the presence of free electrons, these solutions act as powerful reducing agents.
- * The reactivity with water increase on descending the group from Li to Cs and $Li < Na < K < Rb < Cs$ due to increase in electropositive character in the same order.
- * Ionic nature, reducing nature, solubility, basic nature increases from LiH to CsH.
- * LiH is thermally more stable




- * With oxygen (Oxides): When alkali metals are heated in air, Li mainly gives Li_2O , sodium mainly gives sodium peroxide (Na_2O_2) and others give super oxides. (KO_2, RbO_2, CsO_2).

- * Peroxides (O_2^{2-}) are the salts of H_2O_2 . Therefore metal peroxides will give H_2O_2 on reaction with water or dil.acids.



- * Concentrated peroxide solutions will give O_2 on reaction with water



- * Na_2O_2 forms octahydrate crystals $Na_2O_2 \cdot 8H_2O$

- * Na_2O_2 is also known as oxone.


- * Even K_2O_2 can be used for the same purpose and it is much better than Na_2O_2

- * Na_2O_2 is used in qualitative analysis in the identification of chromium salts as it forms yellow colored chromate salts.



- * Super oxides are coloured and paramagnetic due to the presence of unpaired electron or odd electron bond.

Resonance structures of superoxide ion (O_2^-)



- * Super Oxides are paramagnetic in nature.
- * LiO_2 And NaO_2 are yellow in colour and are highly unstable.
- * KO_2 And CsO_2 are orange but RbO_2 is brown coloured.

* Li_2O_2 is white, Na_2O_2 and Cs_2O_2 are yellow coloured. Rb_2O_2 is dark brown, K_2O_2 is orange coloured.

* The monoxides of Li, Na and K are colour less. But the monoxides of Rb and Cs are coloured.

Rb_2O - Yellow: Cs_2O - Orange.

* The solubility of the hydroxides increases from $LiOH$ to $CsOH$

* Cesium halides have body centered cubic lattice.

* Li^+ has maximum degree of hydration and for this reason lithium salts are mostly hydrated

E.g. $LiCl \cdot 2 H_2O$

* The solubilities of bicarbonates in water is less than that of the carbonates.

* $NaHCO_3$ is sparingly soluble in water while Na_2CO_3 is water soluble.

* Except Li_2CO_3 the other carbonates are stable and they decomposing only at very high temperatures. $Li_2CO_3 \rightarrow Li_2O + CO_2$

* The stability increases from Li_2CO_3 to Cs_2CO_3

* Thermal stability of alkali metal carbonates increases from Li_2CO_3 to Cs_2CO_3 . It is because of an increase in the size of cation.

* In alkali metals Li exhibit anomalous behavior. Due to

1. Exceptionally small atomic and ionic size and
2. High polarizing power (i.e. charge/radius ration)
3. Lack of presence of vacant d-orbitals

* Lithium shows diagonal relationship with magnesium.

* Alkyl lithium's are similar to Grignard reagents

Sodium: It is the most abundant element in 1-A group.

* It is used in the preparation of compounds like Na_2O_2 , $NaNH_2$, $NaCN$ etc.

* Na-Pb alloy is used in the preparation of tetraethyl lead (TEL) which is used as anti-knocking agent in petrol.

Potassium (K)

* Na^+ and K^+ play important role in Biology.

* 20% NaOH is formed in the central compartment.

Reactions during electrolysis

Ionisation of brine: $NaCl \rightarrow Na^+ + Cl^-$

At graphite anode: $2Cl^- \rightarrow Cl_2 + 2e^-$

At Hg cathode: $Na^+ + e^- + Hg \rightarrow Na - Hg$

At Hg anode: $Na - Hg \rightarrow Na^+ + e^- + Hg$

At Fe cathode: $2Na^+ + 2e^- + 2H_2O \rightarrow 2NaOH + H_2$

Instead of H_2 gas, sodium is formed in the outer compartments because the discharge potential of sodium is lowered in presence of Hg cathode.

Sodium Carbonate

- * Decahydrate sodium carbonate ($Na_2CO_3 \cdot 10H_2O$) is called washing soda or salt soda.
- * Anhydrous sodium carbonate is called soda ash or soda.
- * It is prepared by
 1. Solvay or ammonia soda process.
- * Solvay process or Ammonia soda process:

Raw materials: Brine, limestone, little NH_3

By-product: $CaCl_2$

Intermediate product: $NaHCO_3$

Recycled products: NH_3 and CO_2



Impurities in Brine solution: Calcium & Magnesium salts.

These are removed in the form of carbonate precipitates.



Precipitation of $NaHCO_3$ in Carbonation tower is due to common ion effect.

Solution from carbonation tower Consists of: $NaHCO_3$ and NH_4Cl
- * It is suitable method to prepare Na_2CO_3 because of low solubility of $NaHCO_3$.
- * K_2CO_3 cannot be manufactured by Solvay's process because $KHCO_3$ is more soluble in water.
- * $Na_2CO_3 \cdot 10H_2O \rightarrow Na_2CO_3 \cdot 7H_2O \rightarrow Na_2CO_3 \cdot H_2O$



- * It's aqueous solution is basic due to hydrolysis of CO_3^{2-}




- * It liberates CO_2 when treated with mineral acids stronger than H_2CO_3 acid.





- * It reacts with sulphur and SO_2 to give hypo



- * When fused with silica it gives sodium silicate or water glass.



- * It causes the precipitation of salt solution as their carbonates.



- * It is used in the manufacture of glass, caustic soda and water glass, Borax, Soap power.

- * It is used for softening of water.

- * It is used in laundries, paper and in dye industry.

- * It is used in sizing of paper.

- * It is used in petroleum refining.


- * It is used in preparation of ultramarine.

- * Sodium bicarbonate or Baking soda ($NaHCO_3$)

- * It is prepared by Solvay's process.

- * It is also obtained by saturating Na_2CO_3 solution with CO_2

- * It is a white crystalline substance preparation



- * $NaHCO_3$ Solution gives yellow colour with methyl orange and it does not give any colour with phenolphthalein.

- * It is used as a fire extinguisher.

- * It is used as an antacid.

www.sakshieducation.com

- * It is used in making baking powders.
- * It is used in the preservation of butter.
- * It is used in the treatment of wool and silk.
- * It is used in effervescent drinks.

www.sakshieducation.com

www.sakshieducation.com