

Electro Chemistry Part- 1

- 1. Which of the following conducts electricity?**
 - 1) Molten Urea
 - 2) Crystalline Sodium Chloride
 - 3) Fused Sodium Chloride
 - 4) Glass
- 2. An electronic conductor in the following is**
 - 1) Solid NaCl
 - 2) Diamond
 - 3) Graphite
 - 4) Aqueous HCl
- 3. The decrease in electrical conductivity of metals with increase in temperature is due to increase in**
 - 1) The velocity of electrons
 - 2) The resistance of the metal
 - 3) The number of electrons
 - 4) The number of metal atoms
- 4. Which of the following is a mixed conductor of electricity?**
 - 1) Aqueous KCl
 - 2) Sodium in liquid NH_3
 - 3) Cane sugar solution
 - 4) CdS & CuS
- 5. In metals and graphite the conduction is due to the flow of**
 - 1) Cations
 - 2) Anions
 - 3) Electrons
 - 4) Both 1&2
- 6. In which of the following, HCl conducts electricity to large extent?**
 - 1) Liquid state
 - 2) In aq. solution
 - 3) In benzene solution.
 - 4) Vapour state
- 7. The reason for increase in electrical conduction of electrolyte with increase in temperature is**
 - A) Increase in the number of ions
 - B) Increase in the speed of ions
 - C) Increase in the degree of dissociation of electrolyte
 - 1) A, B only
 - 2) B, C only
 - 3) A, C only
 - 4) A, B, C

8. Dissociation of an electrolyte in water into negative and positive ions is called

- 1) Electrolysis 2) Hydrolysis 3) Decomposition 4) Ionisation

9. Choose the wrong statement.

- 1) Electrical conductance of an electrolytic conductor increases with increase in temperature.
2) Electrical conductance of a metallic conductor increases with increase in temperature.
3) Electrical conductance of a metallic conductor decreases with increase in temperature.
4) Degree of dissociation of an electrolyte increases with dilution.

10. List - 1

- A) Electronic conductor
B) Non-electrolyte
C) Electrolytic dissociation
D) Arrhenius

List - 2

- 1) Aqueous urea solution
2) Solid sodium
3) Electrolytic conductor
4) Radioactivity increases
5) Conductivity rises with temperature

The correct match is

- | | A | B | C | D | | A | B | C | D |
|----|---|---|---|---|----|---|---|---|---|
| 1) | 5 | 1 | 2 | 3 | 2) | 5 | 2 | 1 | 4 |
| 3) | 2 | 1 | 5 | 3 | 4) | 2 | 5 | 1 | 4 |

11. Which of the following is 100% ionised at any dilution?

- 1) CH_3COOH 2) HCN 3) NaCl 4) NH_4OH

12. Which of the following (1M) conducts more electricity?

- 1) Sulphuric Acid 2) Boric Acid 3) Acetic Acid 4) Phosphorous Acid

13. The degree of dissociation of an electrolyte in aqueous solution depends on

- A) Temperature
B) Concentration of the electrolyte

C) Nature of the electrolyte

- 1) Only A
- 2) Only A, B
- 3) Only B, C
- 4) A, B, C

14. What happens at infinite dilution in a given solution?

- 1) The degree of dissociation is unity for weak electrolytes.
- 2) The electrolyte is 100% ionized.
- 3) All inter ionic attractions disappear.
- 4) All of these.

15. At infinite dilution the degree of dissociation for Urea in aqueous solution is

- 1) 0
- 2) 0.5
- 3) 0.99
- 4) 1

Hint: urea is a non-electrolyte.

16. Choose the correct statement regarding electrolytic cell

- 1) It is a device in which chemical energy is converted into electrical energy.
- 2) Anode is shown by negative sign.
- 3) Oxidation reaction takes place at the anode.
- 4) Electrons flow from cathode to anode.

17. The following are some statements about electrolytic cell.

- A) In this chemical energy is converted into electrical energy.
- B) In this cell electrons flow from anode to cathode
- C) In this cell reduction takes place at cathode
- D) In this cathode is a +ve electrode

The correct combination is


- 1) Only B
- 2) Only C
- 3) Only C, D
- 4) Only B, C

18. The conduction of a salt solution in water depends on the

- 1) Extent of its ionization
- 2) Size of its molecules
- 3) Shape of molecules
- 4) Size of solvent molecules

19. **In the electrolytic cell, flow of electrons is from**
- 1) Cathode to anode in the solution
 - 2) Cathode to anode through external circuit
 - 3) Anode to cathode through external circuit
 - 4) All of these
20. **The unit of specific conductivity is**
- 1) ohms cm^{-1}
 - 2) ohms cm^{-2}
 - 3) ohm $^{-1}$ cm
 - 4) ohm $^{-1}$ cm^{-1}
21. **The unit of equivalent conductivity is**
- 1) ohm cm
 - 2) ohm $^{-1}$ cm^2 (g equivalent) $^{-1}$
 - 3) ohm cm^2 (g equivalent)
 - 4) S cm^{-2}
22. **The equivalent conductance of 1N solution of an electrolyte is nearly**
- 1) Same as its specific conductance
 - 2) 10^{-3} times its specific conductance
 - 3) 10^2 times more than its specific conductance
 - 4) 10^3 times more than its specific conductance
23. **(A): The molar conductance of weak electrolytes is low as compared to that of strong electrolytes at moderate concentrations.**
- (R): Weak electrolytes at moderate concentrations dissociate to a much greater extent when compared to strong electrolytes.**
- 1) Both A and R are true and R is correct explanation to A.
 - 2) Both A and R are true but R is not correct explanation to A.
 - 3) A is true and R is false.
 - 4) Both A and R are false.
24. **Which of the following has highest electrical conductivity in aqueous solutions?**
- 1) 0.1 M acetic acid
 - 2) 0.1 M chloroacetic acid
 - 3) 0.1 M chloroacetic acid
 - 4) 0.1 M tri chloroacetic acid

25. If the specific conductance and conductance of a solution is same, then its cell constant is equal to
 1) 1 2) 0 3) 10 4) 100
26. In electrolysis of dilute H_2SO_4 , what is liberated at anode in presence of inert electrode?
 1) H_2 2) SO_2 3) SO_3 4) O_2
27. Which process occurs in the electrolysis of aqueous solution of nickel chloride at nickel anode?
 1) $\text{Ni}^{2+} + 2\text{e}^- \rightarrow \text{Ni}$ 2) $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$
 3) $2\text{Cl}^- \rightarrow \text{Cl}_2 + 2\text{e}^-$ 4) $\text{Ni} \rightarrow \text{Ni}^{2+} + 2\text{e}^-$
28. Molten CuCl_2 is electrolysed using platinum electrode. The reaction occurring at anode is
 1) $2\text{Cl}^- \rightarrow \text{Cl}_2 (\text{g}) + 2\text{e}^-$ 2) $\text{Cl}_2 (\text{g}) + 2\text{e}^- \rightarrow 2\text{Cl}^-$
 3) $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu} (\text{s})$ 4) $\text{Cu} (\text{s}) \rightarrow \text{Cu}^{2+} + 2\text{e}^-$
29. During the electrolytic reduction of alumina, the reaction at cathode is
 1) $2\text{H}_2\text{O} \rightarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$ 2) $2\text{F}^- \rightarrow \text{F}_2 + 3\text{e}^-$
 3) $\text{Al}^{3+} + 3\text{e}^- \rightarrow \text{Al}$ 4) $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$
30. Specific conductivity of a solution
 1) Increases with dilution 2) Decreases with dilution
 3) Remains unchanged with dilution 4) Depends on mass of electrolyte.
31. A solution of concentration 'C' g equiv/litre has a specific resistance R. The equivalent conductance of the solution is"
 1) R/C 2) C/R 3) $\frac{1000}{\text{RXC}}$ 4) $\frac{1000\text{R}}{\text{C}}$
32. A graph is drawn between the Λ_{eq} values and concentrations of an electrolyte. Which of the following electrolyte will correspond to the graph given?



- 1) KCl 2) CaCl₂ 3) NiSO₄ 4) CH₃COOH

33. For which case 'Λ' values v/s \sqrt{c} shows a straight line

- 1) HCl 2) HCOOH 3) H₃ BO₃ 4) CH₃COOH

Hint: strong electrolytes give straight line.

34. When an aqueous solution of copper sulphate is electrolysed using copper electrodes the reaction at the anode is represented by

- 1) $H^+ + e^- \rightarrow H$ 2) $Cu^{2+} + 2e^- \rightarrow Cu$
 3) $SO_4^{2-}(aq) \rightarrow SO_4 + 2e^-$ 4) $Cu(s) \rightarrow Cu^{2+}(aq) + 2e^-$

35. Aqueous solution of CuSO₄ is electrolysed using inert electrodes till the blue coloured solution becomes colourless. The colourless solution formed is

- 1) Cu (OH)₂ 2) H₂SO₄ 3) CuSO₄ 4) H₂O

36. After the electrolysis of aqueous solution of NaCl using Pt electrodes, the pH of the solution

- 1) Increases 2) Decreases 3) Remains constant 4) Becomes zero

Hint: during electrolysis NaOH is formed.

37. Aqueous solution of AgNO₃ is electrolysed using inert electrodes. At the end of electrolysis

- 1) pH of the solution increases 2) pH of the solution decreases
 3) pH of the solution remains unchanged 4) pH of the solution becomes 14

Hint: during electrolysis HNO₃ is formed.

- 38) 1M aqueous CuSO_4 solution is electrolysed by using copper electrodes for 30 minutes. The concentration of CuSO_4 after electrolysis is

1) 1M 2) 0.75M 3) 0.5M 4) 0.25M

Hint: during electrolysis using active electrodes the composition of electrolyte remains same.

39. According to Kohlrausch law, the limiting value of molar conductivity of an electrolyte A_2B is

1) $\lambda_{\text{A}^+}^\infty + \lambda_{\text{B}^{2-}}^\infty$ 2) $\frac{1}{2}\lambda_{\text{A}^+}^\infty + \lambda_{\text{B}^{2-}}^\infty$

3) $2\lambda_{\text{A}^+}^\infty + \frac{1}{2}\lambda_{\text{B}^{2-}}^\infty$ 4) $2\lambda_{\text{A}^+}^\infty + \lambda_{\text{B}^{2-}}^\infty$

40. The expression showing the relationship between equivalent conductivity and molar conductivity of aq. H_2SO_4 is

1) $\lambda_m = 2 \times \lambda_{\text{eq}}$ 2) $\lambda_{\text{eq}} = 2 \times \lambda_m$ 3) $\lambda_m = 2 / \lambda_{\text{eq}}$ 4) $\lambda_m = 4 \times \lambda_{\text{eq}}$

41. The molar conductivities Λ_{NaOAc}^0 and Λ_{HCl}^0 at infinite dilution in water at 25°C and 91.0 and $426.2 \text{ S cm}^2 / \text{mol}$ respectively. To calculate Λ_{HOAc}^0 the additional value required is

1) Λ_{NaCl}^0 2) $\Lambda_{\text{H}_2\text{O}}^0$ 3) Λ_{NaOH}^0 4) Λ_{KCl}^0

42. The conductivity of 0.001 M acetic acid is $5 \times 10^{-5} \text{ S cm}^{-1}$ and Λ^0 is $500 \text{ S cm}^2 \text{ mol}^{-1}$ then the calculated value of dissociation constant of acetic acid would be

1) 10^{-4} 2) 10^{-5} 3) 10^{-6} 4) 10^{-3}

Solution: $\Lambda_c = K \times 1000 / M = 1000 \times 5 \times 10^{-5} / 0.001 = 50$

Degree of dissociation $\alpha = \Lambda_c / \Lambda_0 = 50 / 500 = 0.1$, $K_a = c\alpha^2 = 0.001 \times (0.1)^2 = 10^{-5}$

43. The distance between two electrodes of a cell is 2.5 cm and area of each electrode is 5 cm² the cell constant (in cm⁻¹) is

- 1) 2 2) 12.5 3) 7.5 4) 0.5

Hint: cell constant=l/a

44. The limiting molar conductivities Λ_0 for NaCl, KBr and KCl are 126, 152 and 150 S. cm² mol⁻¹ respectively. Then Λ_0 for NaBr is

- 1) 128 S cm² mol⁻¹ 2) 302 S cm² mol⁻¹
3) 278 S cm² mol⁻¹ 4) 176 S cm² mol⁻¹

Hint; Λ_0 of NaBr = Λ_0 NaCl + Λ_0 KBr - Λ_0 KCl

45. Which of the following solutions of NaCl has the higher specific conductance?

- 1) 0.001N 2) 0.01N 3) 0.1 N 4) 1 N

Hint: the value of K increases with increase in concentration.

46. Molar conductivity of a solution is 1.26×10^2 S cm² mol⁻¹. Its molarity is 0.01M. Its specific conductivity will be

- 1) 1.26×10^{-5} 2) 1.26×10^{-3} 3) 1.26×10^{-4} 4) 0.0063

Hint; $\lambda_c = K \times 1000/M$

47. The values of equivalent conductivity at infinite dilutions for NH₄Cl, NaOH and NaCl are respectively 149.74, 248.1 and 126.4 ohm⁻¹ cm² equi⁻¹. The value of Λ_{eq} of NH₄OH is

- 1) 371.44 2) 271.44 3) 71.44 4) 224.76

Hint; Λ_{eq} of NH₄OH = Λ_{eq} of NH₄Cl + Λ_{eq} of NaOH + Λ_{eq} of NaCl

48. Molar ionic conductivities of a bivalent electrolyte are 57 and 73. The molar conductivity of the solution will be

- 1) 130 S cm² mol⁻¹ 2) 65 S cm² mol⁻¹ 3) 260 S cm² mol⁻¹ 4) 187 S cm² mol⁻¹

Hint: molar conductivity of the solution = Sum of Molar ionic conductivities = 57 + 73 = 130

49. At a certain temperature and at infinite dilution, the equivalent conductance's of sodium benzoate, hydrochloric acid and sodium chloride are 240, 349 and 229 $\text{ohm}^{-1} \text{cm}^2 \text{equiv}^{-1}$ respectively. The equivalent conductance of benzoic acid in $\text{ohm}^{-1} \text{cm}^2 \text{equiv}^{-1}$ at the same conditions is
- 1) 80 2) 328 3) 360 4) 408

Hint; Λ_o of $\text{C}_6\text{H}_5\text{COOH} = \Lambda_o$ of $\text{C}_6\text{H}_5\text{COONa} + \Lambda_o$ of $\text{HCl} - \Lambda_o$ of NaCl

- 50) The resistance of 1N solution of acetic acid is 250 Ohm. If the cell constant is 1.15 cm^{-1} , then the equivalent conductance will be

- 1) $4.6 \text{ Ohm}^{-1} \text{cm}^2 \text{eq}^{-1}$ 2) $9.2 \text{ Ohm}^{-1} \text{cm}^2 \text{eq}^{-1}$
 3) $18.4 \text{ Ohm}^{-1} \text{cm}^2 \text{eq}^{-1}$ 4) $0.023 \text{ Ohm}^{-1} \text{cm}^2 \text{eq}^{-1}$

Hint: $K = (1/R) \times l/A = 1.15/250 = 4.6 \times 10^{-3}$, $\Lambda = K \times 1000/N = 4.6$

51. The equivalent conductance of 1 M H_2SO_4 solution having conductivity

$26 \times 10^{-2} \text{ ohm}^{-1} \text{cm}^{-1}$ is (in $\text{ohm}^{-1} \text{cm}^2 \text{eq}^{-1}$)

- 1) 260 2) 130 3) 5 4) 10

Hint: for H_2SO_4 Normality $N = MX_2 = 2N$

$$\Lambda_{\text{eq}} = K \times 1000/N = 26 \times 10^{-2} \times 1000/2 = 130$$

52. Equivalent conductance of A_xB_y at infinite dilution will be

- 1) $\lambda^\infty = x\lambda_A^\infty + \lambda_B^\infty$ 2) $\lambda^\infty = x\lambda_A^\infty + y\lambda_B^\infty$
 3) $\lambda^\infty = x\lambda_A^{\infty+} + y\lambda_B^\infty$ 4) All are correct

53. Specific conductivity of 0.1 M solution of KCl at 18°C is 1.12 S. m^{-1} and resistance is 50 ohm. Then cell constant is

- 1) 56 m^{-1} 2) 5.6 m^{-1} 3) 11.2 m^{-1} 4) 1.12 m^{-1}

Hint: cell constant $[l/a] = K \times R$

54. Resistance of 1.0 M aq. solution of an electrolyte is 40 ohm. If area of the electrode of the cell is 3.0 cm^2 & the distance between the electrodes is 1.5 cm, the molar conductivity of the solution is

- 1) $52 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ 2) $24 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 3) $12.5 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ 4) $5.2 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$

55. Equivalent conductance at infinite dilution of BaCl_2 , H_2SO_4 and $\text{HCl}_{(\text{aq})}$ solutions are x_1 , x_2 & x_3 respectively. The equivalent conductance of Ba_2SO_4 at infinite dilution is

- 1) $x_1 + x_2 - 2x_3$ 2) $x_1 + x_2 - x_3$ 3) $x_1 - x_2 + x_3$ 4) $x_1 + 2x_2 + x_3$

56. Equivalent conductance of 1 M CH_3COOH is $10 \text{ ohm}^{-1} \text{ cm}^2 \text{ equiv}^{-1}$ and at infinite dilution is $200 \text{ ohm}^{-1} \text{ cm}^2 \text{ equiv}^{-1}$. The percentage ionisation of CH_3COOH in the 1 M solution is

- 1) 5 % 2) 2% 3) 4% 4) 1%

Hint; $\% \alpha = (\Lambda_c / \Lambda_o) \times 100 = (10/200) \times 100 = 5\%$

57. The specific conductance of 0.1 M HNO_3 is $6.3 \times 10^{-2} \text{ ohm}^{-1} \text{ cm}^{-1}$. The molar conductance of the solution is

- 1) $630 \text{ ohm}^{-1} \text{ cm}^2$ 2) $315 \text{ ohm}^{-1} \text{ cm}^2$ 3) $100 \text{ ohm}^{-1} \text{ cm}^2$ 4) $6300 \text{ ohm}^{-1} \text{ cm}^2$

Hint; $\lambda = K \times 1000 / M$

58. The resistance of 0.01N solution of an electrolyte AB at 328 K is 100 ohm. The specific conductance of solution is (cell constant = 1 cm^{-1})

- 1) 100 ohm 2) 10^{-2} ohm^{-1} 3) 10^2 ohm-cm 4) $10^{-2} \text{ ohm}^{-1} \text{ cm}^{-1}$

Hint; $K = \frac{1}{R} \times \frac{1}{a} = (1/100) \times 1 = 10^{-2} \text{ ohm}^{-1} \text{ cm}^{-1}$

59. For an electrolytic solution of 0.05 mol L^{-1} , the conductivity has been found to be 0.011 S cm^{-1} . The molar conductivity is

- 1) $0.055 \text{ S cm}^2 \text{ mol}^{-1}$ 2) $550 \text{ S cm}^2 \text{ mol}^{-1}$
3) $0.22 \text{ S cm}^2 \text{ mol}^{-1}$ 4) $220 \text{ S cm}^2 \text{ mol}^{-1}$

Hint; $\lambda = \kappa \times 1000 / M$

60. For which of the following electrolyte the value of molar conductivity and equivalent conductivity are same

- 1) Na_2SO_4 2) KCl 3) $\text{Al}_2(\text{SO}_4)_3$ 4) BaCl_2

Key

- 1) 3 2) 3 3) 2 4) 2 5) 3 6) 2 7) 4 8) 4 9) 2 10) 3
11) 3 12) 1 13) 4 14) 4 15) 1 16) 3 17) 4 18) 1 19) 3 20) 4
21) 2 22) 4 23) 3 24) 4 25) 1 26) 4 27) 4 28) 1 29) 3 30) 2
31) 4 32) 4 33) 1 34) 4 35) 2 36) 1 37) 2 38) 1 39) 4
40) 1 41) 1 42) 3 43) 4 44) 1 45) 4 46) 2 47) 2 48) 1 49) 3
50) 1 51) 2 52) 3 53) 1 54) 3 55) 1 56) 1 57) 1 58) 4 59) 4 60) 2