

Spectra, Hydrogen Spectrum, Bohr's Model

- 1. The energy of an electron present in Bohr's second orbit of hydrogen atom is (E - 2001)**
1) $-1312 \text{ J atom}^{-1}$ 2) -328 kJ mol^{-1}
3) -328 J mol^{-1} 4) -164 kJ mol^{-1}
- 2. Splitting of spectral lines under the influence of strong magnetic field is called (AFMC)**
1) Stark Effect 2) Zeeman effect
3) Photoelectric effect 4) None of these
- 3. The wave number of first line of Balmer series of H-atom is 15200 cm^{-1} . The wave number of first Balmer line of Li^{2+} ion is (IIT 92)**
1) 15200 cm^{-1} 2) 60800 cm^{-1}
3) 76000 cm^{-1} 4) $136,800 \text{ cm}^{-1}$
- 4. What are the values of n_1 and n_2 respectively for line in the Lyman series of hydrogen atomic spectrum? (E 2001)**
1) 3 and 5 2) 2 and 3 3) 1 and 3 4) 1 and 4
- 5. If the electron of a hydrogen atom is present in the first orbit, the total energy of the electron is (E-2003)**
1) $\frac{-e^2}{r}$ 2) $\frac{-e^2}{r^2}$ 3) $\frac{-e^2}{2r}$ 4) $\frac{-e^2}{2r^2}$
- 6. The angular momentum of an electron present in the excited state of hydrogen is $1.5h$. The electron is present in (M- 2006)**
1) Third orbit 2) Second orbit
3) Fourth orbit 4) Fifth orbit

7. The wavelength of a spectral line emitted by hydrogen atom in the Lyman series is $16/15R$ cm. What is the value of n_2 ?

(R = Rydberg constant)

(E – 2007)

- 1) 2 2) 3 3) 4 4) 1

8. What is the lowest energy of the spectral line emitted by the hydrogen atom in the Lyman series? (h = Planck constant, c = Velocity of light, R = Rydberg constant)

(M- 2005)

- 1) $\frac{5hcR}{36}$ 2) $\frac{4hcR}{3}$ 3) $\frac{3hcR}{4}$ 4) $\frac{7hcR}{144}$

9. Bohr's radius for the Hydrogen atom ($n=1$) is 0.53Å . The radius for the first excited state is

(CBSE1998)

- 1) 0.13Å 2) 1.06Å 3) 4.77Å 4) 2.12Å

- 10) The wavelength of visible light is

(AIIMS 1998)

- 1) $2000-3700\text{Å}$ 2) $7800-8900\text{Å}$
3) $3800-7600\text{Å}$ 4) $500-1200\text{Å}$

11. The Velocity of the electron in the 2nd orbit of Hydrogen atom is

(AIIMS 2001)

- 1) 10.96×10^6 m/sec 2) 18.88×10^6 m/sec
3) 1.888×10^6 m/sec 4) 1.094×10^6 m/sec

12. In Hydrogen atom energy of the electron in first excited state is -3.4eV .

Then kinetic energy in same orbit is

(CBSE 2002)

- 1) $+3.4\text{eV}$ 2) $+6.8\text{eV}$ 3) -13.6eV 4) $+13.6\text{eV}$

13. The ratio of Radius of 4th and 2nd orbits of H- atom is

(BHU 2003)

- 1) 2 2) 4 3) 3 4) 6

14. When the electron in hydrogen atom jumps from 4th orbit into the first orbit, the frequency of radiation will be

(CBSE 2004)

- 1) 1.54×10^{15} 2) 1.03×10^{15} 3) 3.08×10^{15} 4) 2.0×10^{15}

15. The radius of first orbit in H-atom is R, and then radius of first orbit in will be (PMT2009)

- 1) R/9 2) R/3 3) 3R 4) 9R

16. In Bohr series of lines of hydrogen spectrum, the third line from the red end corresponds to which one of the following inter-orbit jumps of the electron for Bohr orbits is an atom of hydrogen. (2002 A.I.E.E.E)

- 1) $5 \rightarrow 2$ 2) $4 \rightarrow 1$ 3) $2 \rightarrow 5$ 4) $3 \rightarrow 2$

17. What are the values of n_1 and n_2 respectively for H_β line in the Lyman series of hydrogen atomic spectrum? (E-2000)

- 1) 3 and 5 2) 2 and 3 3) 1 and 3 4) 2 and 4

18. What is the wavelength of H_β line the Balmer series of hydrogen spectrum? R = Rydberg constant) (M-2000)

- 1) $36/5R$ 2) $5R/36$ 3) $3R/16$ 4) $16/3R$

KEY

- 1) 2 2) 2 3) 4 4) 4 5) 3 6) 1 7) 3 8) 3

- 9) 4 10) 3 11) 4 12) 1 13) 2 14) 3 15) 2 16) 1

- 17) 3 18) 1

Hints

$$1. E_n = \frac{-1312}{n^2} = \frac{-1312}{2^2} \text{ KJ / mole}$$

$$3. \text{ for same spectral line, } \frac{\bar{\nu}_H}{\bar{\nu}_{Li^{+2}}} = \frac{Z^2_H}{Z^2_{Li^{+2}}} = \frac{1^2}{3^2}$$

$$6. mvr = \frac{nh}{2\pi} = \frac{1.5h}{\pi}, n=3$$

$$7. \frac{1}{\lambda} = R\left(\frac{1}{n_1^2} - \frac{1}{n_2^2}\right), R\left(\frac{1}{1^2} - \frac{1}{n^2}\right) = \frac{15R}{16} \quad n=4$$

8. Lowest energy is for

$$H_{\alpha}, \frac{1}{\lambda} = R\left(\frac{1}{n_1^2} - \frac{1}{n_2^2}\right) = R\left(\frac{1}{1^2} - \frac{1}{2^2}\right) = \frac{3R}{4}$$

$$\Delta E = \frac{hc}{\lambda} = hc\left(\frac{3R}{4}\right) = \frac{3hcR}{4}$$

$$14. \frac{1}{\lambda} = R\left(\frac{1}{n_1^2} - \frac{1}{n_2^2}\right), \nu = \frac{c}{\lambda}$$