GATE 2018 Microbiology (XL-S)

XL(S): Q. 1 – Q. 10 carry one mark each & Q. 11 – Q. 20 carry two marks each.

Q.1	ices in							
	(B) the pathways(C) the modes of	ptors used by viruses required to synthesize v transmission of viruses proteins on the surface of						
Q.2	Which of the follo	Which of the following immune system components can function as an opsonin?						
	(A) Antibodies(C) Histamines		(B) T-cell receptor (D) Interferons	rs				
Q.3	The oral polio vac	ccine (OPV) consists of						
	(A) live attenuate (C) viral toxin	d virus	(B) killed virus (D) viral capsid su	bunit				
Q.4	Which of the following eukaryotic cellular components carries out intracellular degradation during autophagy?							
	(A) Nucleus	(B) Golgi bodies	(C) Ribosomes	(D) Lysosomes				
Q.5	Analysis of DNA sequences suggest that eukaryotic mitochondrial genomes primarily originated from							
	(A) fungi	(B) protozoa	(C) algae	(D) bacteria				
Q.6	Binomial nomeno	elature has NOT yet been	n adopted for					
	(A) bacteria	(B) fungi	(C) viruses	(D) protozoa				
Q.7	Which of the follo	owing is NOT an accept	ted method for steriliz	eation?				
	(A) Autoclaving (C) Gamma rays		(B) X-rays (D) UV rays					
Q.8	The primary product of nitrogen fixation is							
	(A) N ₂	(B) NH ₄ ⁺	(C) NO ₂ ⁻	(D) NO ₃				
Q.9	In humans, the key stages in the life cycle of malarial parasites occur in							
	 (A) red blood cells and the liver (B) red blood cells and platelets (C) red blood cells and the pancreas (D) red blood cells and the gut 							

XL-S 1/3

GATE 2018 Microbiology (XL-S)

Q.10 You have a 50 mg/mL stock solution of arginine. To prepare 1 liter of growth medium for an arginine auxotroph that requires 70 μ g/mL of arginine, the volume of this stock solution that should be added is ______ mL (up to 1 decimal point) .

- Q.11 Accumulating evidence suggest that Domain Archaea is more closely related to Domain Eukarya than to Domain Bacteria. Which of the following properties are shared between eukaryotes and archaea?
 - (i) Protein biogenesis
 - (ii) Presence of sterol containing membranes
 - (iii) Ribosomal subunit structures
 - (iv) Adaptation to extreme environmental conditions
 - (v) Fatty acids with ester linkages in the cell membrane
 - (A) (ii), (iii) and (v)

(B) (i), (ii), (iv), and (v)

(C) (i) and (iii)

(D) (iii) and (iv)

Q.12 Match the antimicrobial agents in group I with their category/mode of action in group II.

Group I		Group II	
(i)	Fluoroquinolones	(p) beta lactam antimicrobial	
(ii)	Amphotericin B	(q) inhibition of protein synthesis	
(iii)	Tetracycline	(r) inhibition of nucleic acid synthesis	
(iv)	Amoxicillin	(s) antifungal agent	

- (A) (i)-(q), (ii)-(s), (iii)-(r), (iv)-(p)
- (B) (i)-(s), (ii)-(r), (iii)-(p), (iv)-(q)
- (C) (i)-(r), (ii)-(s), (iii)-(q), (iv)-(p)
- (D) (i)-(s), (ii)-(r), (iii)-(q), (iv)-(p)
- Q.13 Match the microorganisms to their predominant modes of transmission.

	Microorganism	Mode of Transmission
(i)	Bordetella pertussis	(p) Vector-borne
(ii)	Dengue virus	(q) Blood-borne
(iii)	Entamoeba histolytica	(r) Droplet infection
(iv)	Hepatitis B virus	(s) Contaminated food

- (A) (i)-(r), (ii)-(p), (iii)-(s), (iv)-(q)
- (B) (i)-(s), (ii)-(q), (iii)-(p), (iv)-(r)
- (C) (i)-(q), (ii)-(p), (iii)-(s), (iv)-(r)
- (D) (i)-(s), (ii)-(r), (iii)-(p), (iv)-(q)
- Q.14 Match the precursors/intermediates with the corresponding metabolic pathways.

Precursor/Intermediates	Metabolic pathway
(i) Inosine monophosphate	(p) L-methionine biosynthesis
(ii) Ornithine	(q) L-tryptophan biosynthesis
(iii) Chorismate	(r) Purine biosynthesis
(iv) Homocysteine	(s) L-arginine biosynthesis

- (A) (i)-(q), (ii)-(r), (iii)-(s), (iv)-(p)
- (B) (i)-(p), (ii)-(r), (iii)-(s), (iv)-(q)
- (C) (i)-(r), (ii)-(p), (iii)-(s), (iv)-(q)
- (D) (i)-(r), (ii)-(s), (iii)-(q), (iv)-(p)

2/3

XL-S


GATE 2018 Microbiology (XL-S)

0.15 Match the scientists to their area of major contribution

	Scientists	Area of major contribution
(i)	Antonie van Leeuwenhoek	(p) Taxonomy
(ii)	Carl Linnaeus	(q) Antimicrobial agents
(iii)	Sir Alexander Fleming	(r) Vaccination
(iv)	Louis Pasteur	(s) Microscopy

- (A) (i)-(s), (ii)-(q), (iii)-(p), (iv)-(r)
- (B) (i)-(s), (ii)-(p), (iii)-(q), (iv)-(r)
- (C) (i)-(p), (ii)-(s), (iii)-(r), (iv)-(q)
- (D) (i)-(q), (ii)-(p), (iii)-(r), (iv)-(s)
- Q.16 Which of the following combinations would improve the resolution of a microscope?
 - Increasing the half aperture angle of the objective lens (i)
 - (ii) Decreasing the wavelength of the illumination source
 - Decreasing the numerical aperture of the objective lens (iii)
 - Decreasing the refractive index of immersion medium (iv)
 - (A) (i) and (ii)
- (B) (ii) and (iii)
- (C) (ii) and (iv)
- (D) (i) and (iii)
- Active transport involves the movement of a biomolecule against a concentration gradient across the cell membrane using metabolic energy. If the extracellular concentration of a biomolecule is 0.005M and its intracellular concentration is 0.5M, the least amount of energy that the cell would need to spend to transport this biomolecule from the outside to the inside of the cell is _____ kcal/mol (up to 2 decimal points). (Temperature T = 298K and universal gas constant R = 1.98 cal/mol·K)

A continuous cell culture being carried out in a stirred tank Q.18 reactor is described in terms of its cell mass concentration X and substrate concentration S. The concentration of the substrate in the sterile feed stream is $S_F = 10$ g/L and yield coefficient $Y_{x/s} = 0.5$. The flow rates of the feed stream and the exit stream are equal (F=5 mL/min) and constant. If the specific growth rate (h⁻¹) $\mu = \frac{0.3 \text{ s}}{(1+\text{s})}$, the steady state concentration of S is _____ g/L (up to 1 decimal point).



- The initial concentration of cells (N₀) growing unrestricted in a culture is 1.0×10^6 cells/mL. If the specific growth rate (µ) of the cells is 0.1 h⁻¹, the time required for the cell concentration to become 1.0×10^8 cells/mL is hours (up to 2 decimal points).
- Q.20 The following stoichiometric equation represents the conversion of glucose to lactic acid in a cell:

Glucose +
$$2Pi + 2ADP \rightarrow 2Lactate + 2ATP + 2H_2O$$

If the free energy of conversion of glucose to lactic acid only is $\Delta G^0 = -47000$ cal/mol, the efficiency of energy transfer is ______ % (up to 1 decimal point). $(\Delta G^0 \text{ for ATP hydrolysis is } -7.3 \text{ kcal/mol})$

END OF THE QUESTION PAPER

XL-S 3/3

Q.No.	Туре	Section	Key/Range	Marks
1	MCQ	GA	А	1
2	MCQ	GA	С	1
3	MCQ	GA	В	1
4	MCQ	GA	В	1
5	MCQ	GA	В	1
6	MCQ	GA	А	2
7	MCQ	GA	D	2
8	MCQ	GA	D	2
9	MCQ	GA	В	2
10	MCQ	GA	С	2
1	MCQ	XL-P	Α	1
2	MCQ	XL-P	D	1
3	MCQ	XL-P	D	1
4	NAT	XL-P	11 to 11	1
5	NAT	XL-P	4 to 4	1
6	MCQ	XL-P	D	2
7	MCQ	XL-P	D	2
8	MCQ	XL-P	Α	2
9	MCQ	XL-P	D	2
10	MCQ	XL-P	А	2
11	MCQ	XL-P	С	2
12	MCQ	XL-P	В	2
13	NAT	XL-P	1.39 to 1.43	2

Q.No.	Туре	Section	Key/Range	Marks
14	NAT	XL-P	7.39 to 7.54	2
15	NAT	XL-P	-13.40 to -13.36	2
1	MCQ	XL-Q	В	1
2	MCQ	XL-Q	А	1
3	MCQ	XL-Q	С	1
4	MCQ	XL-Q	С	1
5	MCQ	XL-Q	D	1
6	MCQ	XL-Q	С	1
7	MCQ	XL-Q	D	1
8	MCQ	XL-Q	В	1
9	NAT	XL-Q	12 to 12	1
10	NAT	XL-Q	50 to 50	1
11	MCQ	XL-Q	А	2
12	MCQ	XL-Q	А	2
13	MCQ	XL-Q	D	2
14	MCQ	XL-Q	В	2
15	MCQ	XL-Q	В	2
16	NAT	XL-Q	512 to 512	2
17	NAT	XL-Q	20 to 20	2
18	NAT	XL-Q	0.8 to 0.8	2
19	NAT	XL-Q	77 to 77	2
20	NAT	XL-Q	-8862 to -8862	2
1	MCQ	XL-R	А	1

Q.No.	Туре	Section	Key/Range	Marks
2	MCQ	XL-R	В	1
3	MCQ	XL-R	С	1
4	MCQ	XL-R	D	1
5	MCQ	XL-R	В	1
6	MCQ	XL-R	С	1
7	MCQ	XL-R	А	1
8	MCQ	XL-R	А	1
9	MCQ	XL-R	В	1
10	NAT	XL-R	28.00 to 31.00	1
11	MCQ	XL-R	D	2
12	MCQ	XL-R	D	2
13	MCQ	XL-R	С	2
14	MCQ	XL-R	В	2
15	MCQ	XL-R	С	2
16	MCQ	XL-R	D	2
17	MCQ	XL-R	Α	2
18	MCQ	XL-R	В	2
19	NAT	XL-R	20.25 to 20.25	2
20	NAT	XL-R	11.00 to 12.00	2
1	MCQ	XL-S	В	1
2	MCQ	XL-S	А	1
3	MCQ	XL-S	А	1
4	MCQ	XL-S	D	1

Q.No.	Туре	Section	Key/Range	Marks
5	MCQ	XL-S	D	1
6	MCQ	XL-S	С	1
7	MCQ	XL-S	D	1
8	MCQ	XL-S	В	1
9	MCQ	XL-S	А	1
10	NAT	XL-S	1.38 to1.42	1
11	MCQ	XL-S	С	2
12	MCQ	XL-S	С	2
13	MCQ	XL-S	Α	2
14	MCQ	XL-S	D	2
15	MCQ	XL-S	В	2
16	MCQ	XL-S	Α	2
17	NAT	XL-S	2.60 to 2.80	2
18	NAT	XL-S	0.5 to 0.5	2
19	NAT	XL-S	45.50 to 46.50	2
20	NAT	XL-S	30.5 to 31.5	2
1	MCQ	XL-T	С	1
2	MCQ	XL-T	В	1
3	MCQ	XL-T	Α	1
4	MCQ	XL-T	D	1
5	MCQ	XL-T	С	1
6	MCQ	XL-T	D	1
7	MCQ	XL-T	В	1

Q.No.	Туре	Section	Key/Range	Marks
8	MCQ	XL-T	А	1
9	MCQ	XL-T	А	1
10	MCQ	XL-T	С	1
11	MCQ	XL-T	В	2
12	MCQ	XL-T	D	2
13	MCQ	XL-T	С	2
14	MCQ	XL-T	С	2
15	MCQ	XL-T	В	2
16	MCQ	XL-T	D	2
17	MCQ	XL-T	С	2
18	MCQ	XL-T	Α	2
19	NAT	XL-T	5270 to 5310	2
20	NAT	XL-T	0.056 to 0.062	2
1	MCQ	XL-U	В	1
2	MCQ	XL-U	Α	1
3	MCQ	XL-U	С	1
4	MCQ	XL-U	Α	1
5	MCQ	XL-U	D	1
6	MCQ	XL-U	D	1
7	NAT	XL-U	1.55 to 1.65	1
8	NAT	XL-U	103.0 to 103.2	1
9	NAT	XL-U	54 to 56	1
10	NAT	XL-U	0 to 0	1

Q.No.	Туре	Section	Key/Range	Marks
11	MCQ	XL-U	В	2
12	MCQ	XL-U	С	2
13	MCQ	XL-U	С	2
14	MCQ	XL-U	А	2
15	MCQ	XL-U	D	2
16	MCQ	XL-U	А	2
17	MCQ	XL-U	В	2
18	NAT	XL-U	9.8 to 10.2	2
19	NAT	XL-U	1.1 to 1.8	2
20	NAT	XL-U	10 to 10	2